

▼ This medicinal product is subject to additional monitoring. This will allow quick identification of new safety information. Healthcare professionals are asked to report any suspected adverse reactions. See section 4.8 for how to report adverse reactions.

1. NAME OF THE MEDICINAL PRODUCT

Enhertu 100 mg powder for concentrate for solution for infusion

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

One vial of powder for concentrate for solution for infusion contains 100 mg of trastuzumab deruxtecan. After reconstitution, one vial of 5 mL solution contains 20 mg/mL of trastuzumab deruxtecan (see section 6.6).

Trastuzumab deruxtecan is an antibody-drug conjugate (ADC) that contains a humanised anti-HER2 IgG1 monoclonal antibody (mAb) with the same amino acid sequence as trastuzumab, produced by mammalian (Chinese Hamster Ovary) cells, covalently linked to DXd, an exatecan derivative and a topoisomerase I inhibitor, via a tetrapeptide-based cleavable linker. Approximately 8 molecules of deruxtecan are attached to each antibody molecule.

Excipient with known effect

Each 100 mg vial contains 1.5 mg of polysorbate 80 (E433).

For the full list of excipients, see section 6.1.

3. PHARMACEUTICAL FORM

Powder for concentrate for solution for infusion.

White to yellowish-white lyophilised powder.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Breast cancer

HER2-positive breast cancer

Enhertu in combination with pertuzumab is indicated for the first-line treatment of adult patients with unresectable or metastatic HER2-positive breast cancer (for biomarker-based patient selection, see section 4.2).

Enhertu as monotherapy is indicated for the treatment of adult patients with unresectable or metastatic HER2-positive breast cancer who have received one or more prior anti-HER2-based regimens.

HER2-low and HER2-ultralow breast cancer

Enhertu as monotherapy is indicated for the treatment of adult patients with unresectable or metastatic

- hormone receptor (HR)-positive, HER2-low or HER2-ultralow breast cancer who have received at least one endocrine therapy in the metastatic setting and who are not considered suitable for endocrine therapy as the next line of treatment (see sections 4.2 and 5.1).

- HER2-low breast cancer who have received prior chemotherapy in the metastatic setting or developed disease recurrence during or within 6 months of completing adjuvant chemotherapy (see section 4.2).

Non-small cell lung cancer (NSCLC)

Enhertu as monotherapy is indicated for the treatment of adult patients with advanced NSCLC whose tumours have an activating HER2 (ERBB2) mutation and who require systemic therapy following platinum-based chemotherapy with or without immunotherapy.

Gastric cancer

Enhertu as monotherapy is indicated for the treatment of adult patients with advanced HER2-positive gastric or gastroesophageal junction (GEJ) adenocarcinoma who have received a prior trastuzumab-based regimen.

Other unresectable or metastatic solid tumours

Enhertu as monotherapy is indicated for the treatment of adult patients with unresectable or metastatic HER2-positive (IHC3+) solid tumours who have received prior treatment and who have no satisfactory treatment options (see sections 4.4 and 5.1; for biomarker-based patient selection, see section 4.2).

4.2 Posology and method of administration

Enhertu should be prescribed by a physician and administered under the supervision of a healthcare professional experienced in the use of anticancer medicinal products. In order to prevent medicinal product errors, it is important to check the vial labels to ensure that the medicinal product being prepared and administered is Enhertu (trastuzumab deruxtecan) and not trastuzumab or trastuzumab emtansine.

Enhertu should not be substituted with trastuzumab or trastuzumab emtansine.

Patient selection

HER2-positive breast cancer

Patients treated with trastuzumab deruxtecan or trastuzumab deruxtecan in combination with pertuzumab for breast cancer should have documented HER2-positive tumour status, defined as a score of 3+ by immunohistochemistry (IHC) or a ratio of ≥ 2.0 by *in situ* hybridization (ISH) or by fluorescence *in situ* hybridization (FISH) assessed by a CE-marked *in vitro* diagnostic (IVD) medical device. If a CE-marked IVD is not available, the HER2 status should be assessed by an alternate validated test.

HER2-low or HER2-ultralow breast cancer

Patients treated with trastuzumab deruxtecan should have documented HER2-low tumour status, defined as a score of IHC 1+ or IHC 2+/ISH-, or HER2-ultralow tumour status, described as IHC 0 with membrane staining (IHC $>0<1+$), as assessed by a CE-marked IVD medical device. If a CE-marked IVD is not available, the HER2 status should be assessed by an alternate validated test (see section 5.1).

NSCLC

Patients treated with trastuzumab deruxtecan for advanced NSCLC should have an activating HER2 (ERBB2) mutation detected by a CE-marked *in vitro* diagnostic (IVD) medical device. If a CE-marked IVD is not available, the HER2 mutation status should be assessed by an alternate validated test.

Gastric cancer

Patients treated with trastuzumab deruxtecan for gastric or gastroesophageal junction cancer should have documented HER2-positive tumour status, defined as a score of 3+ by immunohistochemistry (IHC) or a ratio of ≥ 2 by *in situ* hybridization (ISH) or by fluorescence *in situ* hybridization (FISH), assessed by a CE-marked *in vitro* diagnostic (IVD) medical device. If a CE-marked IVD is not available, the HER2 status should be assessed by an alternate validated test.

Other unresectable or metastatic solid tumours

Patients treated with trastuzumab deruxtecan for solid tumours should have documented HER2-positive (IHC 3+) tumour status, assessed by a CE-marked *in vitro* diagnostic (IVD) medical device. If a CE-marked IVD is not available, the HER2 status should be assessed by an alternate validated test.

Posology

Breast cancer, NSCLC, and other unresectable or metastatic solid tumours

The recommended dose of Enhertu is 5.4 mg/kg body weight given as an intravenous infusion once every 3 weeks (21-day cycle) until disease progression or unacceptable toxicity.

Enhertu in combination with pertuzumab (HER2-positive breast cancer)

The recommended dose of Enhertu, in combination with pertuzumab, is 5.4 mg/kg body weight given as an intravenous infusion once every 3 weeks (21-day cycle), with approximately 30 minutes between infusions, until disease progression or unacceptable toxicity.

- Cycle 1, Day 1: Administer 5.4 mg/kg of Enhertu followed by a separate intravenous infusion of pertuzumab at 840 mg.
- Subsequent cycles, Day 1: Administer 5.4 mg/kg of Enhertu followed by a separate intravenous infusion of pertuzumab at 420 mg.

Gastric cancer

The recommended dose of Enhertu is 6.4 mg/kg body weight given as an intravenous infusion once every 3 weeks (21-day cycle) until disease progression or unacceptable toxicity.

The initial dose should be administered as a 90-minute intravenous infusion. If the prior infusion was well tolerated, subsequent doses of Enhertu may be administered as 30-minute infusions.

The infusion rate of Enhertu should be slowed or interrupted if the patient develops infusion-related symptoms (see section 4.8). Enhertu should be permanently discontinued in case of severe infusion reactions.

Premedication

Enhertu is emetogenic (see section 4.8), which includes delayed nausea and/or vomiting. Prior to each dose of Enhertu, patients should be premedicated with a combination regimen of two or three medicinal products (e.g., dexamethasone with either a 5-HT₃ receptor antagonist and/or an NK1 receptor antagonist, as well as other medicinal products as indicated) for prevention of chemotherapy-induced nausea and vomiting.

Dose modifications

Management of adverse reactions may require temporary interruption, dose reduction, or treatment discontinuation of Enhertu per guidelines provided in Tables 1 and 2.

Enhertu dose should not be re-escalated after a dose reduction is made.

Enhertu in combination with pertuzumab

When withholding, modifying, or discontinuing Enhertu, refer to Table 2. Pertuzumab dose modifications should be managed according to the pertuzumab SmPC. Pertuzumab is not to be administered as a single agent.

Table 1: Dose reduction schedule

Dose reduction schedule	Breast cancer, NSCLC, and Other Solid Tumours	Gastric cancer
Recommended starting dose	5.4 mg/kg	6.4 mg/kg
First dose reduction	4.4 mg/kg	5.4 mg/kg
Second dose reduction	3.2 mg/kg	4.4 mg/kg
Requirement for further dose reduction	Discontinue treatment	Discontinue treatment

Table 2: Dose modifications for adverse reactions

Adverse reaction	Severity		Treatment modification
Interstitial lung disease (ILD)/pneumonitis	Asymptomatic ILD/pneumonitis (Grade 1)		Interrupt Enhertu until resolved to Grade 0, then: - if resolved in 28 days or less from date of onset, maintain dose. - if resolved in greater than 28 days from date of onset, reduce dose one level (see Table 1). - consider corticosteroid treatment as soon as ILD/pneumonitis is suspected (see section 4.4).
	Symptomatic ILD/pneumonitis (Grade 2 or greater)		- Permanently discontinue Enhertu. - Promptly initiate corticosteroid treatment as soon as ILD/pneumonitis is suspected (see section 4.4).
Neutropenia	Grade 3 (less than $1.0-0.5 \times 10^9/L$)		Interrupt Enhertu until resolved to Grade 2 or less, then maintain dose.
	Grade 4 (less than $0.5 \times 10^9/L$)		- Interrupt Enhertu until resolved to Grade 2 or less. - Reduce dose by one level (see Table 1).
Febrile neutropenia	Absolute neutrophil count of less than $1.0 \times 10^9/L$ and temperature greater than 38.3 °C or a sustained temperature of 38 °C or greater for more than one hour.		- Interrupt Enhertu until resolved. - Reduce dose by one level (see Table 1).
Left ventricular dysfunction (LVD)	Left ventricular ejection fraction (LVEF) greater than 45% and absolute decrease from baseline is 10% to 20%		Continue treatment with Enhertu.
	LVEF 40% to 45%	And absolute decrease from baseline is less than 10%	- Continue treatment with Enhertu. - Repeat LVEF assessment within 3 weeks.
		And absolute decrease from baseline is 10% to 20%	- Interrupt Enhertu. - Repeat LVEF assessment within 3 weeks. - If LVEF has not recovered to within 10% from baseline, permanently discontinue Enhertu. - If LVEF recovers to within 10% from baseline, resume treatment with Enhertu at the same dose.

Adverse reaction	Severity	Treatment modification
	LVEF less than 40% or absolute decrease from baseline is greater than 20%	• Interrupt Enhertu. • Repeat LVEF assessment within 3 weeks. • If LVEF of less than 40% or absolute decrease from baseline of greater than 20% is confirmed, permanently discontinue Enhertu.
	Symptomatic congestive heart failure (CHF)	• Permanently discontinue Enhertu.

Toxicity grades are in accordance with National Cancer Institute Common Terminology Criteria for Adverse Events Version 5.0 (NCI-CTCAE v.5.0).

Delayed or missed dose

If a planned dose is delayed or missed, it should be administered as soon as possible without waiting until the next planned cycle. The schedule of administration should be adjusted to maintain a 3-week interval between doses. The infusion should be administered at the dose and rate the patient tolerated in the most recent infusion.

Special populations

Elderly

No dose adjustment of Enhertu is required in patients aged 65 years or older. Limited data are available in patients ≥ 75 years of age.

Renal impairment

No dose adjustment is required in patients with mild (creatinine clearance [CL_{Cr}] ≥ 60 and < 90 mL/min) or moderate (CL_{Cr} ≥ 30 and < 60 mL/min) renal impairment (see section 5.2). The potential need for dose adjustment in patients with severe renal impairment or end-stage renal disease cannot be determined as severe renal impairment was an exclusion criterion in clinical studies. A higher incidence of Grade 1 and 2 ILD/pneumonitis leading to an increase in discontinuation of therapy has been observed in patients with moderate renal impairment. In patients with moderate renal impairment at baseline who received Enhertu 6.4 mg/kg, a higher incidence of serious adverse reactions was observed compared to those with normal renal function. Patients with moderate or severe renal impairment should be monitored carefully for adverse reactions including ILD/pneumonitis (see section 4.4).

Hepatic impairment

No dose adjustment is required in patients with total bilirubin ≤ 1.5 times upper limit of normal (ULN), irrespective of aspartate transaminase (AST) value. The potential need for dose adjustment in patients with total bilirubin > 1.5 times ULN, irrespective of AST value, cannot be determined due to limited data; therefore, these patients should be monitored carefully (see sections 4.4 and 5.2).

Paediatric population

The safety and efficacy of Enhertu in children and adolescents below the age of 18 years have not been established. No data are available.

Method of administration

Enhertu is for intravenous use. It must be reconstituted and diluted by a healthcare professional and administered as an intravenous infusion. Enhertu must not be administered as an intravenous push or bolus.

For instructions on reconstitution and dilution of the medicinal product before administration, see section 6.6.

4.3 Contraindications

Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.

4.4 Special warnings and precautions for use

In order to prevent medicinal product errors, it is important to check the vial labels to ensure that the medicinal product being prepared and administered is Enhertu (trastuzumab deruxtecan) and not trastuzumab or trastuzumab emtansine.

Traceability

In order to improve the traceability of biological medicinal products, the name and the batch number of the administered product should be clearly recorded.

Interstitial lung disease/pneumonitis

Cases of interstitial lung disease (ILD), and/or pneumonitis, have been reported with Enhertu or Enhertu in combination with pertuzumab (see section 4.8). Fatal outcomes have been observed. Patients should be advised to immediately report cough, dyspnoea, fever and/or any new or worsening respiratory symptoms. Patients should be monitored for signs and symptoms of ILD/pneumonitis. Evidence of ILD/pneumonitis should be promptly investigated. Patients with suspected ILD/pneumonitis should be evaluated by radiographic imaging, preferably a computed tomography (CT) scan. Consultation with a pulmonologist should be considered. For asymptomatic (Grade 1) ILD/pneumonitis, consider corticosteroid treatment (e.g., ≥ 0.5 mg/kg/day prednisolone or equivalent). Enhertu should be withheld until recovery to Grade 0 and may be resumed according to instructions in Table 2 (see section 4.2). For symptomatic ILD/pneumonitis (Grade 2 or greater), promptly initiate corticosteroid treatment (e.g., ≥ 1 mg/kg/day prednisolone or equivalent) and continue for at least 14 days followed by gradual taper for at least 4 weeks. Enhertu should be permanently discontinued in patients who are diagnosed with symptomatic (Grade 2 or greater) ILD/pneumonitis (see section 4.2). Patients with a history of ILD/pneumonitis or patients with moderate or severe renal impairment may be at increased risk of developing ILD/pneumonitis and should be monitored carefully (see section 4.2). Fatal cases of pneumonia, including two cases adjudicated as ILD, were observed.

Neutropenia

Cases of neutropenia, including febrile neutropenia with a fatal outcome, were reported in clinical studies of Enhertu or Enhertu in combination with pertuzumab. Complete blood counts should be monitored prior to initiation of Enhertu and prior to each dose, and as clinically indicated. Based on the severity of neutropenia, Enhertu may require dose interruption or reduction (see section 4.2).

Left ventricular dysfunction

Left ventricular dysfunction (LVD) has been observed with anti-HER2 therapies. Standard cardiac function testing (echocardiogram or MUGA [multigated acquisition] scanning) should be performed to assess LVEF prior to initiation of Enhertu or Enhertu in combination with pertuzumab and at regular intervals during treatment as clinically indicated. LVD should be managed through treatment interruption. Enhertu should be permanently discontinued if LVEF of less than 40% or absolute decrease from baseline of greater than 20% is confirmed. Enhertu should be permanently discontinued in patients with symptomatic congestive heart failure (CHF) (see Table 2 in section 4.2).

Embryo-foetal toxicity

Enhertu can cause foetal harm when administered to a pregnant woman. In post-marketing reports, use of trastuzumab, a HER2 receptor antagonist, during pregnancy resulted in cases of oligohydramnios manifesting as fatal pulmonary hypoplasia, skeletal abnormalities and neonatal death. Based on

findings in animals and its mechanism of action, the topoisomerase I inhibitor component of Enhertu, DXd, can also cause embryo-foetal harm when administered to a pregnant woman (see section 4.6).

The pregnancy status of females of reproductive potential should be verified prior to the initiation of Enhertu. The patient should be informed of the potential risks to the foetus. Females of reproductive potential should be advised to use effective contraception during treatment and for at least 7 months following the last dose of Enhertu. Male patients with female partners of reproductive potential should be advised to use effective contraception during treatment with Enhertu and for at least 4 months after the last dose of Enhertu (see section 4.6).

Patients with moderate or severe hepatic impairment

There are limited data in patients with moderate hepatic impairment and no data in patients with severe hepatic impairment. As metabolism and biliary excretion are the primary routes of elimination of the topoisomerase I inhibitor, DXd, Enhertu should be administered with caution in patients with moderate and severe hepatic impairment (see sections 4.2 and 5.2).

Efficacy across other unresectable or metastatic solid tumours

The efficacy of Enhertu in other unresectable or metastatic HER2-positive (IHC3+) solid tumours has been established in studies encompassing adult patients (N = 192). Favourable effects of Enhertu in this indication have been shown on the basis of objective response rate and response duration across tumour types. The effect may be quantitatively different depending on tumour type (see section 5.1). For these reasons, Enhertu should only be used if there are no treatment options for which clinical benefit has been established, or where such treatment options have been exhausted (i.e., no satisfactory treatment options).

4.5 Interaction with other medicinal products and other forms of interaction

Co-administration with ritonavir, an inhibitor of OATP1B, CYP3A and P-gp, or with itraconazole, a strong inhibitor of CYP3A and P-gp, resulted in no clinically meaningful (approximately 10-20%) increase in exposures of trastuzumab deruxtecan or the released topoisomerase I inhibitor, DXd. No dose adjustment is required during co-administration of trastuzumab deruxtecan with medicinal products that are inhibitors of CYP3A or OATP1B or P-gp transporters (see section 5.2).

4.6 Fertility, pregnancy and lactation

Women of childbearing potential/Contraception in males and females

Pregnancy status of women of childbearing potential should be verified prior to initiation of Enhertu.

Women of childbearing potential should use effective contraception during treatment with Enhertu and for at least 7 months following the last dose.

Men with female partners of childbearing potential should use effective contraception during treatment with Enhertu and for at least 4 months following the last dose.

Pregnancy

There is no available data on the use of Enhertu in pregnant women. However, trastuzumab, a HER2 receptor antagonist, can cause foetal harm when administered to a pregnant woman. In post-marketing reports, use of trastuzumab during pregnancy resulted in cases of oligohydramnios in some cases manifested as fatal pulmonary hypoplasia, skeletal abnormalities and neonatal death. Based on findings in animals and its mechanism of action, the topoisomerase I inhibitor component of Enhertu, DXd, can be expected to cause embryo-foetal harm when administered to a pregnant woman (see section 5.3).

Administration of Enhertu to pregnant women is not recommended, and patients should be informed of the potential risks to the foetus before they become pregnant. Women who become pregnant must immediately contact their doctor. If a woman becomes pregnant during treatment with Enhertu or within 7 months following the last dose of Enhertu, close monitoring is recommended.

Breast-feeding

It is not known if trastuzumab deruxtecan is excreted in human milk. Human IgG is secreted in human milk, and the potential for absorption and serious adverse reactions to the infant is unknown. Therefore, women should not breast-feed during treatment with Enhertu or for 7 months after the last dose. A decision should be made to discontinue breast-feeding or to discontinue treatment taking into account the benefit of breast-feeding for the child and/or benefit of treatment with Enhertu for the mother.

Fertility

No dedicated fertility studies have been conducted with trastuzumab deruxtecan. Based on results from animal toxicity studies, Enhertu may impair male reproductive function and fertility. It is not known whether trastuzumab deruxtecan or its metabolites are found in seminal fluid. Before starting treatment, male patients should be advised to seek counselling on sperm storage. Male patients must not freeze or donate sperm throughout the treatment period, and for at least 4 months after the final dose of Enhertu.

4.7 Effects on ability to drive and use machines

Enhertu may have a minor influence on the ability to drive and use machines. Patients should be advised to use caution when driving or operating machinery in case they experience fatigue, headache or dizziness during treatment with Enhertu (see section 4.8).

4.8 Undesirable effects

Summary of the safety profile

Enhertu as monotherapy

Enhertu 5.4 mg/kg

The pooled safety population has been evaluated for patients who received at least one dose of Enhertu 5.4 mg/kg (n = 3057) across multiple tumour types in clinical studies. The median duration of treatment in this pool was 9.9 months (range: 0.2 to 76.0 months).

The most common adverse reactions were nausea (70.3%), fatigue (56.9%), vomiting (36.5%), neutropenia (35.9%), anaemia (35.4%), alopecia (34.4%), constipation (33.0%), diarrhoea (30.3%), decreased appetite (29.3%), transaminases increased (27.2%), musculoskeletal pain (24.0%), thrombocytopenia (23.4%) and leukopenia (20.8%).

The most common National Cancer Institute – Common Terminology Criteria for Adverse Events (NCI-CTCAE v.5.0) Grade 3 or 4 adverse reactions were neutropenia (18.5%), anaemia (9.9%), fatigue (8.2%), leukopenia (5.8%), thrombocytopenia (5.2%), nausea (4.8%), lymphopenia (4.2%), transaminases increased (3.7%), hypokalaemia (3.6%), diarrhoea (2.5%), vomiting (2.4%), decreased appetite (1.6%), pneumonia (1.4%) and ejection fraction decreased (1.0%). Grade 5 adverse reactions occurred in 1.1% of patients, including ILD/pneumonitis (1.0%).

Dose interruptions due to adverse reactions occurred in 34.3% of patients treated with Enhertu. The most frequent adverse reactions associated with dose interruption were neutropenia (12.7%), fatigue (4.4%), anaemia (4.3%), upper respiratory tract infection (3.8%), leukopenia (3.1%), ILD/pneumonitis (3.0%), thrombocytopenia (2.8%) and pneumonia (2.5%). Dose reductions occurred in 21.1% of patients treated with Enhertu. The most frequent adverse reactions associated with dose reduction were

fatigue (5.9%), nausea (4.9%), neutropenia (3.3%) and thrombocytopenia (2.2%). Discontinuation of therapy due to an adverse reaction occurred in 10.8% of patients treated with Enhertu. The most frequent adverse reaction associated with permanent discontinuation was ILD/pneumonitis (8.7%).

Enhertu 6.4 mg/kg

The pooled safety population has been evaluated for patients who received at least one dose of Enhertu 6.4 mg/kg (n = 1133), across multiple tumour types in clinical studies. The median duration of treatment in this pool was 5.1 months (range: 0.4 to 41.0 months).

The most common adverse reactions were nausea (64.3%), fatigue (57.3%), anaemia (47.9%), decreased appetite (46.8%), neutropenia (45.9%), vomiting (34.7%), diarrhoea (33.0%), thrombocytopenia (32.9%), leukopenia (31.2%), alopecia (29.0%), constipation (28.2%) and transaminases increased (26.4%).

The most common National Cancer Institute – Common Terminology Criteria for Adverse Events Grade 3 or 4 adverse reactions were neutropenia (28.4%), anaemia (22.8%), leukopenia (12.3%), thrombocytopenia (10.8%), fatigue (8.6%), hypokalaemia (5.8%), pancytopenia (5.6%), nausea (5.6%), lymphopenia (5.5%), decreased appetite (5.3%), transaminases increased (3.6%), pneumonia (3.0%), febrile neutropenia (2.6%), vomiting (2.6%), diarrhoea (1.9%), weight decreased (1.7%), abdominal pain (1.5%), blood alkaline phosphatase increased (1.2%), blood bilirubin increased (1.2%), interstitial lung disease (ILD, 1.1%) and ejection fraction decreased (1.1%). Grade 5 adverse reactions occurred in 2.2% of patients, including ILD (1.6%).

Dose interruptions due to adverse reactions occurred in 40.7% of patients treated with Enhertu. The most frequent adverse reactions associated with dose interruption were neutropenia (14.7%), anaemia (8.5%), fatigue (6.0%), ILD (4.7%), leukopenia (3.9%), pneumonia (3.3%), thrombocytopenia (3.2%), decreased appetite (2.7%) and upper respiratory tract infection (2.6%). Dose reductions occurred in 29.1% of patients treated with Enhertu. The most frequent adverse reactions associated with dose reduction were fatigue (8.4%), neutropenia (6.4%), nausea (5.6%), decreased appetite (4.1%) and thrombocytopenia (3.8%). Discontinuation of therapy due to an adverse reaction occurred in 13.8 % of patients treated with Enhertu. The most frequent adverse reaction associated with permanent discontinuation was ILD (10.1%).

In patients with gastric cancer treated with Enhertu 6.4 mg/kg (n = 546), 19.2% received a transfusion within 28 days after onset of anaemia or thrombocytopenia. Transfusions were primarily for anaemia.

Enhertu 5.4 mg/kg in combination with pertuzumab

The pooled safety population described has been evaluated for patients who received at least one dose of Enhertu 5.4 mg/kg in combination with pertuzumab in a pooled analysis of 431 patients with unresectable or metastatic breast cancer in DESTINY-Breast07 (n = 50) and DESTINY-Breast09 (n = 381). The median duration of treatment was 22 months (range 0.3 to 44.5).

The most common adverse reactions were nausea (74.2%), diarrhoea (64.0%), fatigue (53.4%), neutropenia (49.0%), alopecia (47.8%), vomiting (45.7%), anaemia (40.4%), transaminases increased (38.7%), upper respiratory tract infection (32.0%), constipation (31.3%), decreased appetite (31.3%), leukopenia (29.7%), decreased weight (28.1%), hypokalaemia (26.7%), thrombocytopenia (25.8%), ejection fraction decrease (25.3%), musculoskeletal pain (23.0%) and abdominal pain (22.3%).

The most common National Cancer Institute – Common Terminology Criteria for Adverse Events (NCI-CTCAE v.5.0) Grade 3 or 4 adverse reactions were neutropenia (24.8%), hypokalaemia (13.2%), anaemia (10.7%), diarrhoea (7.4%), fatigue (7.2%), thrombocytopenia (7.0%), transaminases increased (5.3%), leukopenia (5.1%), nausea (4.9%), lymphopenia (3.9%), ejection fraction decreased (3.2%), decreased weight (2.8%), febrile neutropenia (2.3%), decreased appetite (2.1%), vomiting (2.1%), upper respiratory tract infection (1.6%), pneumonia (1.2%) and stomatitis (1.2%).

Grade 5 adverse reactions occurred in 1.6% of patients, including pneumonia (0.9%), ILD/pneumonitis (0.5%), dyspnoea (0.2%) and febrile neutropenia (0.2%).

Dose interruptions due to adverse reactions occurred in 54.1% of patients treated with Enhertu 5.4 mg/kg in combination with pertuzumab. The most frequent adverse reactions (>2%) associated with dose interruption were neutropenia (15.5%), upper respiratory tract infection (12.3%), anaemia (7.2%), fatigue (7.2%), ILD (7.0%), hypokalaemia (6.3%), diarrhoea (5.8%), thrombocytopenia (5.3%), pneumonia (4.6%), leukopenia (3.9%), increased transaminases (3.7%), cough (2.6%), increased blood bilirubin (2.3%), decreased appetite (2.1%), pyrexia (2.1%) and vomiting (2.1%). Dose reductions occurred in 38.5% of patients treated with Enhertu 5.4 mg/kg in combination with pertuzumab. The most frequent adverse reactions (>2%) associated with dose reduction were fatigue (11.6%), neutropenia (7.4%), nausea (6.5%), diarrhoea (6.0%), ILD (3.2%), thrombocytopenia (3.2%), vomiting (3.0%), increased transaminases (2.8%), decreased weight (2.3%) and febrile neutropenia (2.1%). The permanent discontinuation of any drug due to an adverse reaction occurred in 15.5% of patients treated with Enhertu 5.4 mg/kg in combination with pertuzumab. The most frequent adverse reaction (>2%) associated with permanent discontinuation was ILD (6.3%).

Tabulated list of adverse reactions

The adverse reactions in patients who received at least one dose of Enhertu or Enhertu in combination with pertuzumab in clinical studies are presented in Table 3. The adverse reactions are listed by MedDRA system organ class (SOC) and categories of frequency. Frequency categories are defined as: very common ($\geq 1/10$), common ($\geq 1/100$ to $< 1/10$), uncommon ($\geq 1/1,000$ to $< 1/100$), rare ($\geq 1/10,000$ to $< 1/1,000$), very rare ($< 1/10,000$), and not known (cannot be estimated from the available data). Within each frequency grouping, adverse reactions are presented in the order of decreasing seriousness.

Table 3: Adverse reactions in patients treated with trastuzumab deruxtecan

System organ class Frequency category	5.4 mg/kg monotherapy Adverse reaction	6.4 mg/kg monotherapy Adverse reaction	5.4 mg/kg + pertuzumab Adverse reaction
Infections and infestations			
Very common	upper respiratory tract infection ^a	upper respiratory tract infection ^a	upper respiratory tract infection ^a
Common	pneumonia	pneumonia	pneumonia
Blood and lymphatic system disorders			
Very common	anaemia ^b , thrombocytopenia ^d , neutropenia ^c , leukopenia ^e	anaemia ^b , neutropenia ^c , thrombocytopenia ^d , leukopenia ^e , lymphopenia ^f	neutropenia ^c , thrombocytopenia ^d , anaemia ^b , leukopenia ^e , lymphopenia ^f
Common	lymphopenia ^f , pancytopenia ^g	febrile neutropenia, pancytopenia ^g	febrile neutropenia, pancytopenia ^g
Uncommon	febrile neutropenia		
Metabolism and nutrition disorders			
Very common	hypokalaemia ^h , decreased appetite	hypokalaemia ^h , decreased appetite	hypokalaemia ^h , decreased appetite
Common	dehydration	dehydration	dehydration

System organ class Frequency category	5.4 mg/kg monotherapy Adverse reaction	6.4 mg/kg monotherapy Adverse reaction	5.4 mg/kg + pertuzumab Adverse reaction
Nervous system disorders			
Very common	headache ⁱ		headache ⁱ , dizziness, dysgeusia
Common	dizziness, dysgeusia	dizziness, headache ⁱ , dysgeusia	
Eye disorders			
Very common			dry eye
Common	dry eye, vision blurred ^j	dry eye, vision blurred ^j	vision blurred ^j
Respiratory, thoracic and mediastinal disorders			
Very common	interstitial lung disease ^k , cough	interstitial lung disease ^k , cough	interstitial lung disease ^k , cough
Common	dyspnoea, epistaxis	dyspnoea, epistaxis	dyspnoea, epistaxis
Gastrointestinal disorders			
Very common	vomiting, nausea, diarrhoea, abdominal pain ^l , constipation, stomatitis ^m , dyspepsia	nausea, vomiting, diarrhoea, constipation, abdominal pain ^l , stomatitis ^m	diarrhoea, vomiting, nausea, stomatitis ^m , abdominal pain ^l , constipation, dyspepsia
Common	gastritis, abdominal distension, flatulence	dyspepsia, abdominal distension, gastritis, flatulence	gastritis, abdominal distension, flatulence
Hepatobiliary disorders			
Very common	transaminases increased ⁿ	transaminases increased ⁿ	transaminases increased ⁿ
Skin and subcutaneous tissue disorders			
Very common	alopecia	alopecia	rash ^o , pruritus, alopecia
Common	rash ^o , skin hyperpigmentation ^p , pruritus	rash ^o , pruritus, skin hyperpigmentation ^p	dry skin, skin hyperpigmentation ^p
Musculoskeletal and connective tissue disorders			
Very common	musculoskeletal pain ^q	musculoskeletal pain ^q	musculoskeletal pain ^q
General disorders and administration site condition			
Very common	fatigue ^r , pyrexia	fatigue ^r , pyrexia, oedema peripheral	fatigue ^r , pyrexia
Common	oedema peripheral		mucosal inflammation

System organ class Frequency category	5.4 mg/kg monotherapy Adverse reaction	6.4 mg/kg monotherapy Adverse reaction	5.4 mg/kg + pertuzumab Adverse reaction
Investigations			
Very common	ejection fraction decreased ^s , weight decreased	ejection fraction decreased ^s , weight decreased	ejection fraction decreased ^s , weight decreased, blood alkaline phosphatase increased
Common	blood bilirubin increased ^t , blood creatinine increased, blood alkaline phosphatase increased	blood alkaline phosphatase increased, blood bilirubin increased ^t , blood creatinine increased	blood bilirubin increased ^t , blood creatinine increased
Injury, poisoning and procedural complications			
Common	infusion-related reactions ^u		infusion-related reactions ^u
Uncommon		infusion-related reactions ^u	

^a For all tumour types at 5.4 mg/kg and all tumour types at 6.4 mg/kg, includes influenza, influenza-like illness, nasopharyngitis, pharyngitis, sinusitis, rhinitis, laryngitis and upper respiratory tract infection. For 5.4 mg/kg + pertuzumab, includes influenza, influenza-like illness, laryngitis, nasopharyngitis, pharyngitis, rhinitis, sinusitis and upper respiratory tract infection.

^b For all tumour types at 5.4 mg/kg, includes anaemia, haemoglobin decreased, red blood cell count decreased and haematocrit decreased. For all tumour types at 6.4 mg/kg, includes anaemia, haemoglobin decreased, haematocrit decreased and red blood cell count decreased. For 5.4 mg/kg + pertuzumab, includes anaemia, haematocrit decreased, haemoglobin decreased and red blood cell count decreased.

^c Includes neutropenia and neutrophil count decreased.

^d For all tumour types at 5.4 mg/kg and all tumour types at 6.4 mg/kg, includes thrombocytopenia and platelet count decreased. For 5.4 mg/kg + pertuzumab, includes platelet count decreased and thrombocytopenia.

^e Includes leukopenia and white blood cell count decreased.

^f For all tumour types at 5.4 mg/kg and all tumour types at 6.4 mg/kg, includes lymphopenia and lymphocyte count decreased. For 5.4 mg/kg + pertuzumab, includes lymphocyte count decreased and lymphopenia.

^g Pancytopenia was defined as a subject that met all 3 criteria of Haemoglobin level < 100 g/L, Neutrophils < 1.5x10⁹/L and Platelets < 100x10⁹/L based on the same lab sample collection date and/or the preferred term pancytopenia.

^h For all tumour types at 5.4 mg/kg and all tumour types at 6.4 mg/kg, includes hypokalaemia and blood potassium decreased. For 5.4 mg/kg + pertuzumab, includes blood potassium decreased and hypokalaemia.

ⁱ For all tumour types at 5.4 mg/kg, includes headache, sinus headache and migraine. For all tumour types at 6.4 mg/kg and 5.4 mg/kg + pertuzumab, includes headache and migraine.

^j Includes vision blurred and visual impairment.

^k For all tumour types at 5.4 mg/kg, interstitial lung disease includes events that were adjudicated ILD: acute respiratory failure (n = 2), alveolitis (n = 2), atypical pneumonia (n = 1), bronchiectasis (n = 1), bronchitis (n = 1), disease progression (n = 1), hypersensitivity pneumonitis (n = 1), idiopathic interstitial pneumonia (n = 1), interstitial lung disease (n = 121), lower respiratory tract infection (n = 1), lung disorder (n = 1), lung infiltration (n = 1), lung opacity (n = 4), lymphangitis (n = 1), organising pneumonia (n = 13), pneumonia (n = 9), pneumonia bacterial (n = 2), pneumonia fungal (n = 1), pneumonitis (n = 160), pulmonary fibrosis (n = 2), pulmonary mass (n = 1), pulmonary toxicity (n = 4), radiation pneumonitis (n = 6) and respiratory failure (n = 5). For all tumour types at 6.4 mg/kg, interstitial lung disease includes events that were adjudicated ILD: alveolitis (n = 1), interstitial lung disease (n = 68), lung opacity (n = 2), organising pneumonia (n = 4), pneumonia (n = 1), pneumonitis (n = 98), pulmonary toxicity (n = 1), radiation pneumonitis (n = 1) and respiratory failure (n = 5). For 5.4 mg/kg + pertuzumab, interstitial lung disease includes events that were adjudicated ILD: chronic obstructive pulmonary disease (n = 1), interstitial lung disease (n = 25), organising pneumonia (n = 1), pneumonia (n = 4), pneumonitis (n = 27) and radiation pneumonitis (n = 2).

- ^l For all tumour types at 5.4 mg/kg and all tumour types at 6.4 mg/kg, includes abdominal discomfort, gastrointestinal pain, abdominal pain, abdominal pain lower and abdominal pain upper. For 5.4 mg/kg + pertuzumab, includes abdominal discomfort, abdominal pain, abdominal pain lower, abdominal pain upper and gastrointestinal pain.
- ^m For all tumour types at 5.4 mg/kg, includes stomatitis, aphthous ulcer, mouth ulceration, oral mucosa erosion and oral mucosal eruption. For all tumour types at 6.4 mg/kg, includes stomatitis, aphthous ulcer and mouth ulceration. For 5.4 mg/kg + pertuzumab, includes aphthous ulcer, mouth ulceration, oral mucosal eruption and stomatitis.
- ⁿ For all tumour types at 5.4 mg/kg and all tumour types at 6.4 mg/kg, includes transaminases increased, alanine aminotransferase increased, aspartate aminotransferase increased, gamma-glutamyltransferase increased, hepatic function abnormal, liver function test abnormal, liver function test increased and hypertransaminasaemia. For 5.4 mg/kg + pertuzumab, includes alanine aminotransferase increased, aspartate aminotransferase increased, gamma-glutamyl transferase increased, hepatic function abnormal, hypertransaminasaemia, liver function test increased and transaminases increased.
- ^o For all tumour types at 5.4 mg/kg, includes rash, rash pustular, rash maculo-papular, rash papular, rash macular and rash pruritic. For all tumour types at 6.4 mg/kg, includes rash, rash pustular, rash maculo-papular, rash papular and rash pruritic. For 5.4 mg/kg + pertuzumab, includes rash, rash macular, rash maculo-papular, rash pruritic and rash pustular.
- ^p For all tumour types at 5.4 mg/kg, includes skin hyperpigmentation, skin discolouration and pigmentation disorder. For all tumour types at 6.4 mg/kg, includes skin hyperpigmentation and pigmentation disorder. For 5.4 mg/kg + pertuzumab, includes pigmentation disorder, skin discolouration and skin hyperpigmentation.
- ^q For all tumour types at 5.4 mg/kg and all tumour types at 6.4 mg/kg, includes back pain, myalgia, pain in extremity, musculoskeletal pain, muscle spasms, bone pain, neck pain, musculoskeletal chest pain and limb discomfort. For 5.4 mg/kg + pertuzumab, includes back pain, bone pain, muscle spasms, musculoskeletal chest pain, musculoskeletal pain, myalgia, neck pain and pain in extremity.
- ^r For all tumour types at 5.4 mg/kg and all tumour types at 6.4 mg/kg, includes asthenia, fatigue, malaise and lethargy. For 5.4 mg/kg + pertuzumab, includes asthenia, fatigue, lethargy and malaise.
- ^s For all tumour types at 5.4 mg/kg, ejection fraction decreased includes laboratory parameters of LVEF decrease (n = 434) and/or preferred terms of ejection fraction decreased (n = 170), cardiac failure (n = 5), cardiac failure acute (n = 1), cardiac failure chronic (n = 2), cardiac failure congestive (n = 1) and left ventricular dysfunction (n = 4). For all tumour types at 6.4 mg/kg, ejection fraction decreased includes laboratory parameters of LVEF decrease (n = 125) and/or preferred terms of ejection fraction decreased (n = 20), left ventricular dysfunction (n = 1), cardiac failure (n = 2), cardiac failure acute (n = 1) and cardiac failure congestive (n = 1). For 5.4 mg/kg + pertuzumab, ejection fraction decreased includes laboratory parameters of LVEF decrease (n = 104) and/or preferred terms of ejection fraction decreased (n = 36), cardiac failure (n = 5), cardiac failure chronic (n = 1) and left ventricular dysfunction (n = 5).
- ^t For all tumour types at 5.4 mg/kg, includes blood bilirubin increased, hyperbilirubinaemia, bilirubin conjugated increased and blood bilirubin unconjugated increased. For all tumour types at 6.4 mg/kg, includes blood bilirubin increased, hyperbilirubinaemia and bilirubin conjugated increased. For 5.4 mg/kg + pertuzumab, includes bilirubin conjugated increased, blood bilirubin increased, blood bilirubin unconjugated increased and hyperbilirubinaemia.
- ^u For all tumour types at 5.4 mg/kg, cases of infusion-related reactions include infusion-related reaction (n = 28) and hypersensitivity (n = 3). For all tumour types at 6.4 mg/kg, cases of infusion-related reactions include infusion-related reaction (n = 6) and hypersensitivity (n = 1). The majority of the cases of infusion-related reactions were Grade 1 and Grade 2. For 5.4 mg/kg + pertuzumab, includes infusion-related reaction (n = 10).

Description of selected adverse reactions

Interstitial lung disease/pneumonitis

Enhertu as monotherapy

In patients treated with Enhertu 5.4 mg/kg in clinical studies across multiple tumour types (n = 2553), ILD, pneumonitis, organising pneumonia and acute interstitial pneumonitis were reported by the investigator in 13.8% of patients. ILD/pneumonitis was confirmed by adjudication in 12.9% of patients, leading to drug discontinuation in 8.7% of patients and drug interruption in 3.0% of patients. Most ILD/pneumonitis cases were Grade 1 (3.0%) and Grade 2 (8.1%). Grade 3 cases occurred in 0.7% and one Grade 4 case occurred. Grade 5 (fatal) events occurred in 1.0% of patients. Median time to first onset was 5.5 months (range: -0.3 to 40.2), including two patients adjudicated as having pre-existing ILD. Recovery was not reported for 27.4% of patients with adjudicated ILD/pneumonitis at a median follow up of 286 days (see sections 4.2 and 4.4).

In patients treated with Enhertu 6.4 mg/kg in clinical studies across multiple tumour types (n = 1133), ILD, pneumonitis, organising pneumonia, and acute interstitial pneumonitis were reported by the investigator in 16.9% of patients. ILD/pneumonitis was confirmed by adjudication in 15.4% of patients, leading to drug discontinuation in 10.1% of patients and drug interruption in 4.7% of patients. Most ILD/pneumonitis cases were Grade 1 (4.1%) and Grade 2 (8.6%). Grade 3 cases occurred in 1.1% and one Grade 4 case occurred. Grade 5 (fatal) events occurred in 1.6% of patients. Median time to first onset was 4.1 months (range: -0.5 to 21.0), including two patients adjudicated as having pre-existing ILD. Recovery was not reported for 37.4% of patients with adjudicated ILD/pneumonitis at a median follow up of 251 days (see sections 4.2 and 4.4).

Enhertu 5.4 mg/kg in combination with pertuzumab

In patients treated with Enhertu 5.4 mg/kg in combination with pertuzumab (n = 431), ILD, pneumonitis, organising pneumonia and acute interstitial pneumonitis were reported by the investigator in 16.7% of patients. ILD was confirmed by adjudication in 13.0% of patients, leading to drug discontinuation in 6.3% of patients and drug interruption in 7.0% of patients. Most ILD cases were Grade 1 (3.9%) and Grade 2 (8.4%). Grade 3 cases occurred in 0.2%. No Grade 4 case occurred. Grade 5 (fatal) events occurred in 0.5% of patients. Median time to first onset was 8.1 months (range: 0.6 to 33.8). Recovery was not reported for 26.8% of patients with adjudicated ILD/pneumonitis at a median follow up of 336 days (see sections 4.2 and 4.4).

Neutropenia

Enhertu as monotherapy

In patients treated with Enhertu 5.4 mg/kg in clinical studies (n = 3057) across multiple tumour types, neutropenia was reported in 35.9% of patients and 18.5% had Grade 3 or 4 events. Median time to onset was 43 days (range: 0 days to 51.5 months), and median duration of the first event was 21 days (range: 1 day to 30.8 months). Febrile neutropenia was reported in 0.9% of patients and one event (<0.1%) was Grade 5 (see section 4.2).

In patients treated with Enhertu 6.4 mg/kg in clinical studies across multiple tumour types (n = 1133), neutropenia was reported in 45.9% of patients and 28.4% had Grade 3 or 4 events. Median time to onset was 16 days (range: 1 day to 24.8 months), and median duration of the first event was 9 days (range: 1 day to 17.2 months). Febrile neutropenia was reported in 2.6% of patients and 0.1% were Grade 5 (see section 4.2).

Enhertu 5.4 mg/kg in combination with pertuzumab

In patients treated with Enhertu 5.4 mg/kg in combination with pertuzumab in clinical studies (n = 431), neutropenia was reported in 49% of patients and 24.8% had Grade 3 or 4 events. Median time to first onset was 42 days (range: 1 day to 36.8 months) and median duration of the first event was 20 days (range: 1 day to 27.6 months). Febrile neutropenia was reported in 2.6% of patients and one event (0.2%) was Grade 5 (see section 4.2).

Left ventricular dysfunction

Enhertu as monotherapy

In patients treated with Enhertu 5.4 mg/kg in clinical studies across multiple tumour types (n = 3057), left ventricular dysfunction was reported in 181 patients (5.9%), of which 17 (0.6%) were Grade 1, 147 (4.8%) were Grade 2, 16 (0.5%) were Grade 3 and 1 (<0.1%) was Grade 4. The observed frequency of LVEF decreased based on laboratory parameters (echocardiogram or MUGA scanning) was 411/2740 (15.0%) for Grade 2 and 22/2740 (0.8%) for Grade 3. Treatment with Enhertu has not been studied in patients with LVEF less than 50% prior to initiation of treatment (see section 4.2).

Left ventricular dysfunction led to treatment interruption in 31/3057 (1.0%) patients. The median time to worst grade LVEF was 6.9 months, and the median time to recovery ($\geq 90\%$ baseline) from worst grade LVEF was 2.8 months.

In patients treated with Enhertu 6.4 mg/kg in clinical studies across multiple tumour types (n = 1133), LVEF decrease was reported in 23 patients (2.0%), of which 1 (0.1%) was Grade 1, 16 (1.4%) were Grade 2, and 6 (0.5%) were Grade 3. The observed frequency of LVEF decreased based on laboratory parameters (echocardiogram or MUGA scanning) was 114/953 (12.0%) for Grade 2, and 11/953 (1.2%) for Grade 3.

Left ventricular dysfunction led to treatment interruption in 6/1133 (0.5%) patients. The median time to worst grade LVEF was 5.5 months, and the median time to recovery ($\geq 90\%$ baseline) from worst grade LVEF was 2.8 months.

Enhertu 5.4 mg/kg in combination with pertuzumab

In patients treated with Enhertu 5.4 mg/kg in combination with pertuzumab (n = 431), left ventricular dysfunction was reported in 46 patients (10.7%), of which 4 (0.9%) were Grade 1, 33 (7.7%) were Grade 2, 8 (1.9%) were Grade 3 and 1 (0.2%) Grade 4. The observed frequency of LVEF decreased based on laboratory parameters (echocardiogram or MUGA scanning) was 90/407 (22.1%) for Grade 2 and 14/407 (3.4%) for Grade 3. Treatment with Enhertu has not been studied in patients with LVEF less than 50% prior to initiation of treatment (see section 4.2).

Left ventricular dysfunction led to treatment discontinuation in 8/431 (1.9%) patients and treatment interruption in 6/431 (1.4%) patients. The median time to worst grade LVEF was 11.6 months and the median time to recovery ($\geq 90\%$ baseline) from worst grade LVEF was 3.0 months.

Infusion-related reactions

Enhertu as monotherapy

In patients treated with Enhertu 5.4 mg/kg in clinical studies (n = 3057) across multiple tumour types, infusion-related reactions were reported in 31 patients (1.0%), the majority of which were Grade 1 or Grade 2 severity. Six events (0.2%) of infusion-related reactions led to dose interruptions, and one event ($<0.1\%$) led to discontinuation.

In patients treated with Enhertu 6.4 mg/kg in clinical studies (n = 1133) across multiple tumour types, infusion-related reactions were reported in 7 patients (0.6%), all of which were Grade 1 or Grade 2 severity. No Grade 3 events were reported. One event (0.1%) of infusion-related reaction led to dose interruption, and no events led to discontinuation.

Enhertu 5.4 mg/kg in combination with pertuzumab

In patients treated with Enhertu 5.4 mg/kg in combination with pertuzumab in clinical studies (n = 431), infusion-related reactions were reported in 10 patients (2.3%), the majority of which were Grade 1 or Grade 2 severity. Seven events (1.6%) of infusion-related reactions led to dose interruptions and no events led to discontinuation.

Immunogenicity

As with all therapeutic proteins, there is a potential for immunogenicity. Across 5.4 mg/kg and 6.4 mg/kg doses evaluated in clinical studies, 2.2% (70/3124) of evaluable patients developed antibodies against trastuzumab deruxtecan following treatment with Enhertu. The incidence of treatment-emergent neutralising antibodies against trastuzumab deruxtecan was 0.1% (3/3124). There was no apparent effect between development of antibodies on the pharmacokinetics, safety and/or effectiveness of Enhertu.

Paediatric population

Safety has not been established in this population.

Elderly

Enhertu as monotherapy

In patients treated with Enhertu 5.4 mg/kg in clinical studies across multiple tumour types (n = 3057), 27.1% were 65 years or older and 5.8% were 75 years or older. There was a higher incidence of Grade 3-4 adverse reactions observed in patients aged 65 years or older (48.2%) as compared to patients younger than 65 years old (44.3%), leading to more discontinuations due to adverse reactions. The incidence of fatal adverse reactions was 2.2% in patients aged 65 years or older and 0.8% in patients younger than 65 years of age.

Of the 1133 patients across multiple tumour types in clinical studies treated with Enhertu 6.4 mg/kg, 39.6% were 65 years or older and 7.9% were 75 years or older. The incidence of Grade 3-4 adverse reactions observed in patients 65 years or older was 60.8% and 61.1% in younger patients. There was a higher incidence of Grade 3-4 adverse reactions observed in patients 75 years of age or older (64.4%) compared to patients less than 75 years of age (60.7%). In patients 75 years or older, there was a higher incidence of serious adverse reactions (34.4%) and fatal events (4.4%) compared to patients less than 75 years (21.2% and 1.6%). Data are limited to establish the safety in patients 75 years or older.

Enhertu 5.4 mg/kg in combination with pertuzumab

In patients treated with Enhertu 5.4 mg/kg in combination with pertuzumab (n = 431), 17.4% were 65 years or older and 3% were 75 years or older. The incidence of Grade 3-4 adverse reactions observed in patients aged 65 years or older was 52% and in younger patients was 55.1%. The incidence of fatal adverse reactions was 4.0% in patients aged 65 years or older and 1.1% in patients younger than 65 years of age. The incidence of adjudicated ILD/pneumonitis was 20.0% in patients aged 65 years or older and 11.5% in patients younger than 65 years of age.

Ethnic differences

In clinical studies, no relevant differences in exposure or efficacy were observed between patients of different ethnic groups. Asian patients receiving Enhertu 6.4 mg/kg had a higher incidence ($\geq 10\%$ difference) of neutropenia (58.3% vs. 29.4%), anaemia (55.2% vs. 38.3%), leukopenia (46.7% vs. 10.5%), and thrombocytopenia (43.1% vs. 19.3%) compared to non-Asian patients. In Asian patients, 3.4% experienced a bleeding event within 14 days after onset of thrombocytopenia compared to 0.8% of non-Asian patients.

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via the HPRA Pharmacovigilance website: www.hpra.ie.

4.9 Overdose

The maximum tolerated dose of trastuzumab deruxtecan has not been determined. In clinical studies, single doses higher than 8.0 mg/kg have not been tested. In case of overdose, patients must be closely monitored for signs or symptoms of adverse reactions and appropriate symptomatic treatment initiated.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antineoplastic agents, HER2 (Human Epidermal Growth Factor Receptor 2) inhibitors, ATC code: L01FD04

Mechanism of action

Enhertu, trastuzumab deruxtecan, is a HER2-targeted antibody-drug conjugate. The antibody is a humanised anti-HER2 IgG1 attached to deruxtecan, a topoisomerase I inhibitor (DXd) bound by a tetrapeptide-based cleavable linker. The antibody-drug conjugate is stable in plasma. The function of the antibody portion is to bind to HER2 expressed on the surface of certain tumour cells. After binding, the trastuzumab deruxtecan complex then undergoes internalisation and intracellular linker cleavage by lysosomal enzymes that are upregulated in cancer cells. Upon release, the membrane-permeable DXd causes DNA damage and apoptotic cell death. DXd, an exatecan derivative, is approximately 10 times more potent than SN-38, the active metabolite of irinotecan.

In vitro studies indicate that the antibody portion of trastuzumab deruxtecan, which has the same amino acid sequence as trastuzumab, also binds to FcγRIIIa and complement C1q. The antibody mediates antibody-dependent cellular cytotoxicity (ADCC) in human breast cancer cells that overexpress HER2. In addition, the antibody inhibits signalling through the phosphatidylinositol 3-kinase (PI3-K) pathway in human breast cancer cells that overexpress HER2.

Clinical efficacy

HER2-positive breast cancer

DESTINY-Breast09 (NCT04784715)

The efficacy and safety of Enhertu in combination with pertuzumab were evaluated in DESTINY-Breast09, a Phase 3, randomised, three-arm, multicentre, global study that enrolled 1157 adult patients with HER2-positive (IHC 3+ or ISH+) advanced, or metastatic breast cancer, as determined at a central laboratory using PATHWAY/VENTANA anti-HER2/neu (4B5) and VENTANA HER2 Dual ISH DNA Probe Cocktail.

The study included adult patients with unresectable or metastatic HER2-positive breast cancer who had not received prior chemotherapy or HER2-targeted therapy for advanced or metastatic breast cancer. Patients were eligible if they (a) had *de novo* metastatic breast cancer disease or (b) had received neoadjuvant or adjuvant chemotherapy and/or HER2-targeted therapy which was completed more than 6 months before the diagnosis of advanced or metastatic disease. A single line of prior endocrine therapy was permitted for advanced or metastatic breast cancer. The study excluded patients with a history of ILD/pneumonitis requiring treatment with steroids or ILD/pneumonitis at screening, patients with symptomatic brain metastases, or ECOG performance status >1 and patients with a history of clinically significant cardiac disease.

Patients were randomised 1:1:1 to receive either Enhertu 5.4 mg/kg monotherapy (n = 387), Enhertu 5.4 mg/kg plus pertuzumab (n = 383), or THP (taxane [docetaxel or paclitaxel], trastuzumab and pertuzumab) (n = 387) by intravenous infusion every three weeks. Patients who received Enhertu plus pertuzumab or THP and who had HR positive tumours (n = 416) were allowed to receive concurrent endocrine therapy after 6 cycles of Enhertu or after discontinuation of the taxane in the THP arm. This occurred in 13.5% (n = 28) of patients with HR positive tumours in the Enhertu plus pertuzumab arm and 38.3% (n = 80) of patients in the THP arm, respectively.

The primary efficacy outcome measure was progression-free survival (PFS) as assessed by blinded independent central review (BICR) based on RECIST v1.1.

Overall survival (OS) was the key secondary efficacy outcome measure. PFS based on investigator assessment, confirmed objective response rate (ORR) and duration of response (DOR) were secondary endpoints.

In the overall population, demographic and baseline disease characteristics were balanced between treatment arms. Of the 1157 patients randomised, the median age was 54 years (range: 20-88); 82% were <65 years; 100% were female; 50% of the patients were Asian, 37.2% were White, and

2.6% were Black or African American. The percentage of patients who had *de novo* disease was 52.2% and recurrent disease was 47.8%. The percentage of patients who had HR positive tumours was 53.2%, HR negative was 46.8% and 30.9% of patients had a PIK3CA mutation.

Efficacy results are summarised in Table 4 and Figure 1. OS data were immature at PFS interim analysis.

Table 4: Efficacy results in DESTINY-Breast09

Efficacy Parameter	Enhertu 5.4 mg/kg + pertuzumab* N = 383	THP [§] N = 387
Progression-Free Survival (PFS) per BICR		
Number of events (%)	118 (30.8)	172 (44.4)
Median, months (95% CI)	40.7 (36.5, NE)	26.9 (21.8, NE)
Hazard ratio (95% CI)	0.56 (0.44, 0.71)	
p-value ^{† ¶}	< 0.00001	
Overall Survival (OS)		
Number of events (%)	58 (15.1)	68 (17.6)
Median, months (95% CI)	NE	NE
Hazard ratio (95% CI)	0.84 (0.59, 1.19)	
Confirmed Objective Response Rate (ORR) per BICR [‡]		
n (%)	326 (85.1)	304 (78.6)
95% CI	(81.2, 88.5)	(74.1, 82.5)

CI = confidence interval; NE = not estimable

[§] THP = taxane [docetaxel or paclitaxel], trastuzumab and pertuzumab.


* Patients who discontinued Enhertu due to toxicity prior to disease progression could continue study treatment with trastuzumab and this occurred in 8.7% of patients in the Enhertu in combination with pertuzumab arm.

[†] At this DCO (26 Feb 2025), interim analysis criterion for superiority (p-value <0.00043) already met for Enhertu in combination with pertuzumab vs THP. Enhertu plus placebo remains blinded until final PFS analysis, which is planned after a predefined number of PFS events across all arms.

[‡] The p-value was calculated using a stratified log-rank test.

[‡] Not multiplicity controlled.

Figure 1: Kaplan-Meier plot of progression-free survival per BICR



DESTINY-Breast03 (NCT03529110)

The efficacy and safety of Enhertu were studied in DESTINY-Breast03, a multicentre, open-label, active-controlled, randomised, two-arm phase 3 study that enrolled patients with HER2-positive, unresectable or metastatic breast cancer who received prior trastuzumab and taxane therapy for metastatic disease or developed disease recurrence during or within 6 months of completing adjuvant therapy.

Archival breast tumour samples were required to show HER2 positivity defined as HER2 IHC 3+ or ISH-positive. The study excluded patients with a history of ILD/pneumonitis requiring treatment with steroids or ILD/pneumonitis at screening, patients with untreated and symptomatic brain metastases, patients with a history of clinically significant cardiac disease and patients with prior treatment with an anti-HER2 antibody-drug conjugate in the metastatic setting. Patients were randomised 1:1 to receive either Enhertu 5.4 mg/kg (N = 261) or trastuzumab emtansine 3.6 mg/kg (N = 263) administered by intravenous infusion once every three weeks. Randomisation was stratified by hormone receptor status, prior treatment with pertuzumab, and history of visceral disease. Treatment was administered until disease progression, death, withdrawal of consent, or unacceptable toxicity.

The primary efficacy outcome measure was progression-free survival (PFS) as evaluated by blinded independent central review (BICR) according to Response Evaluation Criteria in Solid Tumours (RECIST v1.1). Overall survival (OS) was a key secondary efficacy outcome measure. PFS based on investigator assessment, confirmed objective response rate (ORR), and duration of response (DOR) were secondary endpoints.

Patient demographics and baseline disease characteristics were balanced between treatment arms. Of the 524 patients randomised, the baseline demographic and disease characteristics were: median age 54 years (range: 20 to 83); 65 years or older (20.2%); female (99.6%); Asian (59.9%), White (27.3%), Black or African American (3.6%); Eastern Cooperative Oncology Group (ECOG) performance status 0 (62.8%) or 1 (36.8%); hormone receptor status (positive: 51.9%); presence of visceral disease (73.3%); presence of brain metastases at baseline (15.6%); and 48.3% of patients received one line of prior systemic therapy in the metastatic setting. The percentage of patients who had not received prior treatment for metastatic disease was 9.5%. The percentage of patients who were previously treated with pertuzumab was 61.1%.

At the prespecified interim analysis for PFS based on 245 events (73% of total events planned for final analysis), the study showed a statistically significant improvement in PFS per BICR in patients randomised to Enhertu compared to trastuzumab emtansine. PFS by BICR data from the primary analysis (data cutoff 21 May 2021) and updated OS, ORR and DOR results from data cutoff 25 July 2022 are presented in Table 5.

Table 5: Efficacy results in DESTINY-Breast03

Efficacy parameter	Enhertu N = 261	trastuzumab emtansine N = 263
Progression-free survival (PFS) per BICR^a		
Number of events (%)	87 (33.3)	158 (60.1)
Median, months (95% CI)	NR (18.5, NE)	6.8 (5.6, 8.2)
Hazard ratio (95% CI)	0.28 (0.22, 0.37)	
p-value	p < 0.000001 [†]	
Overall survival (OS)^b		
Number of events (%)	72 (27.6)	97 (36.9)
Median, months (95% CI)	NR (40.5, NE)	NR (34.0, NE)
Hazard ratio (95% CI)	0.64 (0.47, 0.87)	
p-value ^c	p = 0.0037	
PFS per BICR (updated)^b		
Number of events (%)	117 (44.8)	171 (65.0)
Median, months (95% CI)	28.8 (22.4, 37.9)	6.8 (5.6, 8.2)
Hazard ratio (95% CI)	0.33 (0.26, 0.43)	
Confirmed objective response rate (ORR) per BICR^b		
n (%)	205 (78.5)	92 (35.0)
95% CI	(73.1, 83.4)	(29.2, 41.1)
Complete response n (%)	55 (21.1)	25 (9.5)
Partial response n (%)	150 (57.5)	67 (25.5)
Duration of response per BICR^b		
Median, months (95% CI)	36.6 (22.4, NE)	23.8 (12.6, 34.7)

CI = confidence interval; NE = not estimable; NR = not reached

[†]Presented as 6 decimal places

^a Data cutoff 21 May 2021

^b Data cutoff 25 July 2022 for a pre-planned OS interim analysis

^c The p-value is based on a stratified log-rank test; crossed the efficacy boundary of 0.013.



Figure 2: Kaplan-Meier plot of overall survival (Data cutoff 25 July 2022)

Figure 3: Kaplan-Meier plot of progression-free survival per BICR (Data cutoff 25 July 2022)



Similar PFS results were observed across prespecified subgroups including prior pertuzumab therapy, hormone receptor status, and presence of visceral disease.

DESTINY-Breast02 (NCT03523585)

The efficacy and safety of Enhertu were evaluated in study DESTINY-Breast02, a Phase 3, randomised, multicentre, open-label, active-controlled study that enrolled patients with unresectable or metastatic HER2-positive breast cancer, who were resistant or refractory to prior T-DM1 therapy. Archival breast tumour samples were required to show HER2 positivity defined as HER2 IHC 3+ or ISH-positive. The study excluded patients with a history of ILD/pneumonitis requiring treatment with steroids or ILD/pneumonitis at screening, patients with untreated and symptomatic brain metastases and patients with a history of clinically significant cardiac disease. Patients were randomised 2:1 to receive either Enhertu 5.4 mg/kg (n = 406) by intravenous infusion every three weeks, or treatment of physician's choice (n = 202, trastuzumab plus capecitabine or lapatinib plus capecitabine). Randomisation was stratified by hormone receptor status, prior treatment with pertuzumab and history of visceral disease. Treatment was administered until disease progression, death, withdrawal of consent or unacceptable toxicity.

The primary efficacy outcome measure was progression-free survival (PFS) as assessed by blinded independent central review (BICR) based on RECIST v1.1. Overall survival (OS) was a key secondary efficacy outcome measure. PFS based on investigator assessment, confirmed objective response rate (ORR) and duration of response (DOR) were secondary objectives.

Demographic and baseline disease characteristics were similar between treatment arms. Of the 608 patients randomised, the median age was 54 years (range 22 to 88); female (99.2%); White (63.2%), Asian (29.3%), Black or African American (2.8%); Eastern Cooperative Oncology Group (ECOG) performance status 0 (57.4%) or 1 (42.4%); hormone receptor status (positive: 58.6%); presence of visceral disease (78.3%); presence of brain metastases at baseline (18.1%) and 4.9% of patients received one line of prior systemic therapy in the metastatic setting.

Efficacy results are summarised in Table 6 and Figures 4 and 5.

Table 6: Efficacy results in DESTINY-Breast02

Efficacy parameter	Enhertu N = 406	Treatment of physician's choice N = 202
PFS per BICR		
Number of events (%)	200 (49.3)	125 (61.9)
Median, months (95% CI)	17.8 (14.3, 20.8)	6.9 (5.5, 8.4)
Hazard ratio (95% CI)	0.36 (0.28, 0.45)	
p-value	p < 0.000001†	
Overall survival (OS)		
Number of events (%)	143 (35.2)	86 (42.6)
Median, months (95% CI)	39.2 (32.7, NE)	26.5 (21.0, NE)
Hazard ratio (95% CI)	0.66 (0.50, 0.86)	
p-value ^a	p = 0.0021	
PFS per investigator assessment		
Number of events (%)	206 (50.7)	152 (75.2)
Median, months (95% CI)	16.7 (14.3, 19.6)	5.5 (4.4, 7.0)
Hazard ratio (95% CI)	0.28 (0.23, 0.35)	
Confirmed objective response rate (ORR) per BICR		
n (%)	283 (69.7)	59 (29.2)
95% CI	(65.0, 74.1)	(23.0, 36.0)
Complete response n (%)	57 (14.0)	10 (5.0)
Partial response n (%)	226 (55.7)	49 (24.3)
Duration of response per BICR		
Median, months (95% CI)	19.6 (15.9, NE)	8.3 (5.8, 9.5)

CI = confidence interval; NE = not estimable

† Presented as 6 decimal places



^a The p-value is based on a stratified log-rank test; crossed the efficacy boundary of 0.004.**Figure 4: Kaplan-Meier plot of progression-free survival per BICR**

Figure 5: Kaplan-Meier plot of overall survival



DESTINY-Breast01 (NCT03248492)

The efficacy and safety of Enhertu were studied in DESTINY-Breast01, a multicentre, open-label, single-arm Phase 2 study that enrolled patients with HER2-positive, unresectable and/or metastatic breast cancer who had received two or more prior anti-HER2-based regimens, including trastuzumab emtansine (100%), trastuzumab (100%) and pertuzumab (65.8%). Archival breast tumour samples were required to show HER2 positivity defined as HER2 IHC 3+ or ISH-positive. The study excluded patients with a history of treated ILD or ILD at screening, patients with untreated or symptomatic brain metastases, and patients with a history of clinically significant cardiac disease. Patients enrolled had at least 1 measurable lesion per RECIST v1.1. Enhertu was administered by intravenous infusion at 5.4 mg/kg once every three weeks until disease progression, death, withdrawal of consent, or unacceptable toxicity. The primary efficacy outcome measure was confirmed objective response rate (ORR) according to RECIST v1.1 in the intent-to-treat (ITT) population as evaluated by independent central review (ICR). The secondary efficacy outcome measure was duration of response (DOR).

Of the 184 patients enrolled in DESTINY-Breast01, baseline demographic and disease characteristics were: median age 55 years (range: 28 to 96); 65 years or older (23.9%); female (100%); White (54.9%), Asian (38.0%), Black or African American (2.2%); Eastern Cooperative Oncology Group (ECOG) performance status 0 (55.4%) or 1 (44.0%); hormone receptor status (positive: 52.7%); presence of visceral disease (91.8%); previously treated and stable brain metastases (13.0%); median number of prior therapies in the metastatic setting: 5 (range: 2 to 17); sum of diameters of target lesions (< 5 cm: 42.4%, ≥ 5 cm: 50.0%).

An earlier analysis (median duration of follow-up 11.1 months [range: 0.7 to 19.9 months]) showed a confirmed objective response rate of 60.9% (95% CI: 53.4, 68.0) with 6.0% being complete responders and 54.9% being partial responders; 36.4% had stable disease, 1.6% had progressive disease and 1.1% were not evaluable. Median duration of response at that time was 14.8 months (95% CI: 13.8, 16.9) with 81.3% of responders having a response of ≥ 6 months (95% CI: 71.9, 87.8). Efficacy results from an updated data cutoff with median duration of follow-up of 20.5 months (range: 0.7 to 31.4 months) are shown in Table 7.

Table 7: Efficacy results in DESTINY-Breast01 (intent-to-treat analysis set)

	DESTINY-Breast01 N = 184
Confirmed objective response rate (95% CI)*†	61.4% (54.0, 68.5)
Complete response (CR)	6.5%
Partial response (PR)	54.9%
Duration of response‡	
Median, months (95% CI)	20.8 (15.0, NR)
% with duration of response ≥ 6 months (95% CI)§	81.5% (72.2, 88.0)

ORR 95% CI calculated using Clopper-Pearson method

CI = confidence interval

95% CIs calculated using Brookmeyer-Crowley method

*Confirmed responses (by blinded independent central review) were defined as a recorded response of either CR/PR, confirmed by repeat imaging not less than 4 weeks after the visit when the response was first observed.

†Of the 184 patients, 35.9% had stable disease, 1.6% had progressive disease and 1.1% were not evaluable.

‡Includes 73 patients with censored data

§Based on Kaplan-Meier estimation

NR = not reached

Consistent anti-tumour activity was observed across prespecified subgroups based on prior pertuzumab therapy and hormone receptor status.

HER2-low and HER2-ultralow breast cancer

DESTINY-Breast06 (NCT04494425)

The efficacy and safety of Enhertu were evaluated in study DESTINY-Breast06, a randomised, multicentre, open-label Phase 3 study that randomised 866 adult patients with advanced or metastatic HR+ breast cancer with HER2-low (IHC 1+ or IHC 2+/ISH-) or HER2-ultralow expression as determined by the PATHWAY/VENTANA anti-HER2/neu (4B5) evaluated at a central laboratory. HER2-ultralow (IHC 0 with membrane staining, described as IHC >0<1+ in the study) is defined as faint and incomplete membrane HER2 staining that is seen in 10% or fewer tumour cells. Patients were eligible if they had disease progression on (a) at least 2 lines of endocrine therapy in the metastatic setting or (b) one line of endocrine therapy in the metastatic setting and demonstrated progression within 24 months of the start of adjuvant endocrine therapy, or within 6 months of starting first line endocrine therapy in combination with a CDK 4/6 inhibitor in the metastatic setting. Patients with prior chemotherapy in the neo-adjuvant or adjuvant setting were eligible if they had a disease-free interval greater than 12 months. The study excluded patients with prior chemotherapy for advanced or metastatic disease, patients with a history of ILD/pneumonitis requiring treatment with steroids or ILD/pneumonitis at screening, uncontrolled or significant cardiovascular disease, untreated and symptomatic brain metastases, or ECOG performance status >1.

Patients were randomised 1:1 to receive either Enhertu 5.4 mg/kg (N = 436) by intravenous infusion every three weeks or physician's choice of single agent chemotherapy (N = 430, capecitabine 60%, nab-paclitaxel 24%, or paclitaxel 16%). Randomisation was stratified by prior CDK4/6 inhibitor use (yes or no), prior taxane use in the non-metastatic setting (yes or no), and HER2 IHC status of tumour samples (IHC 2+/ISH-, IHC 1+, IHC >0 <1+). Treatment with Enhertu was administered until disease progression, death, withdrawal of consent, or unacceptable toxicity.

The primary efficacy outcome measure was PFS in patients with HER2-low breast cancer assessed by BICR based on RECIST v1.1. Key secondary efficacy outcome measures were PFS assessed by BICR based on RECIST v1.1 in the overall population (HER2-low and HER2-ultralow), OS in HER2-low patients, and OS in the overall population. ORR and DOR were secondary endpoints.

In the overall population, demographics and baseline tumour characteristics were similar between treatment arms. Of the 866 patients randomised, the median age was 57 years (range: 28 to 87); 31% were age 65 or older; 99.9% were female; 53% were White, 35% were Asian, and 1% were Black or

African American. Patients had an ECOG performance status of 0 (59%) or 1 (39%) at baseline; 18% were IHC >0<1+, 55% were IHC 1+, 27% were IHC 2+/ISH-; 67% had liver metastases, 32% had lung metastases, 8% had brain metastases, and 3% had bone-only metastases. Patients had a median of 2 prior lines of endocrine therapy in the metastatic setting (range: 1 to 5) with 17% having 1 and 68% having 2. Eighty-nine percent of patients had prior endocrine therapy in combination with CDK4/6i treatment in the metastatic setting, 47% had prior anthracycline use, and 41% had prior taxane use in the non-metastatic setting.

Efficacy results are summarised in Table 8 and Figures 6 and 7.

Table 8: Efficacy Results in DESTINY-Breast06

Table 8. Efficacy Results in DESTINY-Breast0				
Efficacy Parameter	HER2-low		Overall Population (HER2-low and HER2-ultralow)	
	Enhertu (N = 359)	Chemotherapy (N = 354)	Enhertu (N = 436)	Chemotherapy (N = 430)
Progression Free Survival per BICR				
Number of events (%)	225 (62.7)	232 (65.5)	269 (61.7)	271 (63.0)
Median, months (95% CI)	13.2 (11.4, 15.2)	8.1 (7.0, 9.0)	13.2 (12.0, 15.2)	8.1 (7.0, 9.0)
Hazard ratio (95% CI)	0.62 (0.52, 0.75)		0.64 (0.54, 0.76)	
p-value	<0.0001		<0.0001	
Overall Survival*				
Number of events (%)	136 (37.9)	146 (41.2)	161 (36.9)	174 (40.5)
Median, months (95% CI)	28.9 (25.7, 33.7)	27.1 (23.5, 29.9)	28.9 (26.4, 32.7)	27.4 (23.9, 29.9)
Hazard ratio (95% CI)	0.83 (0.66, 1.05)		0.81 (0.66, 1.01)	
Confirmed Objective Response Rate per BICR†				
n (%)	203 (56.5)	114 (32.2)	250 (57.3)	134 (31.2)
95% CI	51.2, 61.7	27.4, 37.3	52.5, 62.0	26.8, 35.8
Duration of Response per BICR‡				
Median, months (95% CI)	14.1 (11.8, 15.9)	8.6 (6.7, 11.3)	14.3 (12.5, 15.9)	8.6 (6.9, 11.5)

Data cutoff: 18 March 2024

CI = confidence interval

*First planned interim analysis

†Results were not controlled for type 1 error and should be interpreted descriptively

Consistent PFS benefit was observed across multiple prespecified subgroups, including HER2 expression (IHC >0 <1+, IHC 1+, IHC 2+/ISH-), prior CDK4/6 inhibitor use (yes or no), prior taxane use in the non-metastatic setting (yes or no), and number of prior lines of endocrine therapy in the metastatic setting.

In the HER2-ultralow subgroup (n = 152), median PFS was 13.2 months (95% CI: 9.8, 17.3) in patients randomised to Enhertu (N = 76) and 8.3 months (95% CI: 5.8, 15.2) in patients randomised to chemotherapy with a hazard ratio of 0.78 (95% CI: 0.50, 1.21). Median OS was 29.5 months (95% CI: 27.9, NE) in patients randomised to Enhertu and 27.4 months (95% CI: 19.4, NE) in patients randomised to chemotherapy with a hazard ratio of 0.75 (95% CI: 0.43, 1.29). Confirmed objective response rate was 61.8% (95% CI: 50.0, 72.8) and 26.3% (95% CI: 16.9, 37.7) in patients randomised to Enhertu and chemotherapy, respectively. Median duration of response was 14.3 months (95% CI: 9.2, 20.7) and 14.1 months (95% CI: 5.9, not estimable) in patients randomised to Enhertu and chemotherapy, respectively.

Figure 6: Kaplan-Meier Plot of Progression Free Survival (Overall Population)





Figure 7: Kaplan-Meier Plot of Overall Survival (Overall Population)



DESTINY-Breast04 (NCT03734029)

The efficacy and safety of Enhertu were studied in DESTINY-Breast04, a phase 3, randomised, multicentre, open-label study that enrolled 557 adult patients with unresectable or metastatic HER2-low breast cancer. The study included 2 cohorts: 494 hormone receptor positive (HR+) patients and 63 hormone receptor negative (HR-) patients. HER2-low expression was defined as IHC 1+ (defined as faint, partial staining of the membrane in greater than 10% of the cancer cells) or IHC 2+/ISH-, as determined by the PATHWAY/VENTANA anti-HER2/neu (4B5) evaluated at a central laboratory. Patients must have received chemotherapy in the metastatic setting or have developed disease recurrence during or within 6 months of completing adjuvant chemotherapy. According to the inclusion criteria, patients who were HR+ must have received at least one endocrine therapy and be ineligible for further endocrine therapy at the time of randomisation. Patients were randomised 2:1 to receive either Enhertu 5.4 mg/kg (N = 373) by intravenous infusion every three weeks or physician's choice of chemotherapy (N = 184, eribulin 51.1%, capecitabine 20.1%, gemcitabine 10.3%, nab paclitaxel 10.3%, or paclitaxel 8.2%). Randomisation was stratified by HER2 IHC status of tumour samples (IHC 1+ or IHC 2+/ISH-), number of prior lines of chemotherapy in the metastatic setting (1 or 2) and HR status/prior CDK4/6i treatment (HR+ with prior CDK4/6 inhibitor treatment, HR+ without prior CDK4/6 inhibitor treatment, or HR-). Treatment was administered until disease progression, death, withdrawal of consent, or unacceptable toxicity. The study excluded patients with a history of ILD/pneumonitis requiring treatment with steroids or ILD/pneumonitis at screening and clinically significant cardiac disease. Patients were also excluded for untreated or symptomatic brain metastases or ECOG performance status > 1.

The primary efficacy endpoint was progression-free survival (PFS) in patients with HR+ breast cancer assessed by BICR based on RECIST v1.1. Key secondary efficacy endpoints were PFS assessed by BICR based on RECIST v1.1 in the overall population (all randomised HR+ and HR- patients), overall survival (OS) in HR+ patients and OS in the overall population. ORR, DOR and patient-reported outcomes (PROs) were secondary endpoints.

Demographics and baseline tumour characteristics were similar between treatment arms. Of the 557 patients randomised, the median age was 57 years (range: 28 to 81); 23.5% were age 65 or older; 99.6% were female and 0.4% were male; 47.9% were White, 40.0% were Asian and 1.8% were Black or African American. Patients had an ECOG performance status of 0 (54.8%) or 1 (45.2%) at baseline; 57.6% were IHC 1+, 42.4% were IHC 2+/ISH-; 88.7% were HR+ and 11.3% HR-; 69.8% had liver metastases, 32.9% had lung metastases, and 5.7% had brain metastases. The percentage of patients who had prior anthracycline use in the (neo)adjuvant setting was 46.3% and 19.4% in the locally advanced and/or metastatic setting. In the metastatic setting, patients had a median of 3 prior lines of systemic therapy (range: 1 to 9) with 57.6% having 1 and 40.9% having 2 prior chemotherapy regimens; 3.9% were early progressors (progression in the neo/adjuvant setting). In HR+ patients, the median number of prior lines of endocrine therapy was 2 (range: 0 to 9) and 70% had prior CDK4/6 inhibitor treatment.

Efficacy results are summarised in Table 9 and Figures 8 and 9.

Table 9: Efficacy results in DESTINY-Breast04

Efficacy parameter	HR+ cohort		Overall population (HR+ and HR- cohort)	
	Enhertu (N = 331)	Chemotherapy (N = 163)	Enhertu (N = 373)	Chemotherapy (N = 184)
Overall survival				
Number of events (%)	126 (38.1)	73 (44.8)	149 (39.9)	90 (48.9)
Median, months (95% CI)	23.9 (20.8, 24.8)	17.5 (15.2, 22.4)	23.4 (20.0, 24.8)	16.8 (14.5, 20.0)

Efficacy parameter	HR+ cohort		Overall population (HR+ and HR- cohort)	
	Enhertu (N = 331)	Chemotherapy (N = 163)	Enhertu (N = 373)	Chemotherapy (N = 184)
Hazard ratio (95% CI)	0.64 (0.48, 0.86)		0.64 (0.49, 0.84)	
p-value	0.0028		0.001	
Progression-free survival per BICR				
Number of events (%)	211 (63.7)	110 (67.5)	243 (65.1)	127 (69.0)
Median, months (95% CI)	10.1 (9.5, 11.5)	5.4 (4.4, 7.1)	9.9 (9.0, 11.3)	5.1 (4.2, 6.8)
Hazard ratio (95% CI)	0.51 (0.40, 0.64)		0.50 (0.40, 0.63)	
p-value	< 0.0001		< 0.0001	
Confirmed objective response rate per BICR*				
n (%)	175 (52.6)	27 (16.3)	195 (52.3)	30 (16.3)
95% CI	47.0, 58.0	11.0, 22.8	47.1, 57.4	11.3, 22.5
Complete Response n (%)	12 (3.6)	1 (0.6)	13 (3.5)	2 (1.1)
Partial Response n (%)	164 (49.2)	26 (15.7)	183 (49.1)	28 (15.2)
Duration of response per BICR*				
Median, months (95% CI)	10.7 (8.5, 13.7)	6.8 (6.5, 9.9)	10.7 (8.5, 13.2)	6.8 (6.0, 9.9)

CI = confidence interval

*Based on data from electronic case report form for the HR+ cohort: N = 333 for Enhertu arm and N = 166 chemotherapy arm.

Consistent OS and PFS benefit were observed across prespecified subgroups, including HR status, prior CDK4/6i treatment, number of prior chemotherapies and IHC 1+ and IHC 2+/ISH- status. In the HR- subgroup, median OS was 18.2 months (95% CI: 13.6, not estimable) in patients randomised to Enhertu compared to 8.3 months (95% CI: 5.6, 20.6) in patients randomised to chemotherapy with a hazard ratio of 0.48 (95% CI: 0.24, 0.95). Median PFS was 8.5 months (95% CI: 4.3, 11.7) in patients randomised to Enhertu and 2.9 months (95% CI: 1.4, 5.1) in patients randomised to chemotherapy with a hazard ratio of 0.46 (95% CI: 0.24, 0.89).

At an updated descriptive analysis with a median follow-up of 32 months, OS improvements were consistent with the primary analysis. The HR in the overall population was 0.69 (95% CI: 0.55, 0.86) with a median OS of 22.9 months (95% CI: 21.2, 24.5) in the Enhertu arm versus 16.8 months (95% CI: 14.1, 19.5) in the chemotherapy arm. The Kaplan-Meier curve for the updated OS analysis is shown in Figure 8.

Figure 8: Kaplan-Meier plot of overall survival (overall population) (updated analysis)





Figure 9: Kaplan-Meier plot of progression-free survival per BICR (overall population)



NSCLC

DESTINY-Lung02 (NCT04644237)

The efficacy and safety of Enhertu were studied in DESTINY-Lung02, a phase 2, randomised study evaluating two dose levels. The treatment dosage assignment was blinded to patients and investigators. The study included adult patients with metastatic HER2-mutant NSCLC who had received at least one regimen containing platinum-based chemotherapy. Identification of an activating HER2 (ERBB2) mutation was prospectively determined in tumour tissue by local laboratories using a validated test such as next generation sequencing, polymerase chain reaction or mass spectrometry. Patients were

randomised 2:1 to receive Enhertu 5.4 mg/kg or 6.4 mg/kg every 3 weeks, respectively. Randomisation was stratified by prior anti-programmed cell death receptor-1 (PD-1) and/or anti-programmed cell death ligand 1 (PD-L1) treatment (yes versus no). Treatment was administered until disease progression, death, withdrawal of consent, or unacceptable toxicity. The study excluded patients with a history of ILD/pneumonitis requiring treatment with steroids or ILD/pneumonitis at screening and clinically significant cardiac disease. Patients were also excluded for untreated and symptomatic brain metastases or ECOG performance status >1.

The primary efficacy outcome measure was confirmed ORR as assessed by BICR using RECIST v1.1. The secondary efficacy outcome measure was DOR.

Demographic and baseline disease characteristics from the 102 patients enrolled in the 5.4 mg/kg arm were: median age 59.4 years (range 31 to 84); female (63.7%); Asian (63.7%), White (22.5%), or Other (13.7%); ECOG performance status 0 (28.4%) or 1 (71.6%); 97.1% had a mutation in the ERBB2 kinase domain, 2.9% in the extracellular domain; 96.1% had a HER2 mutation in exon 19 or exon 20; 34.3% had stable brain metastases; 46.1% were former smokers, none were current smokers; 21.6% had a prior lung resection. In the metastatic setting, 32.4% had greater than 2 prior systemic therapies, 100% received platinum-based therapy, 73.5% received anti-PD-1/PD-L1 therapy, and 50.0% had prior treatment with platinum therapy and anti-PD-1/PD-L1 therapy in combination.

Efficacy results are summarised in Table 10. The median duration of follow-up was 11.5 months (data cutoff: 23 December 2022).

Table 10: Efficacy results in DESTINY-Lung02

Efficacy parameter	DESTINY-Lung02 5.4 mg/kg N = 102
Confirmed objective response rate (ORR) per BICR	
n (%)	50 (49.0)
(95% CI)*	(39.0, 59.1)
Complete response (CR) n (%)	1 (1.0)
Partial response (PR) n (%)	49 (48.0)
Duration of response	
Median, months (95% CI)†	16.8 (6.4, NE)

*95% CI calculated using Clopper-Pearson method

CI = confidence interval, NE = not estimable

†95% CI calculated using Brookmeyer-Crowley method

Gastric cancer

DESTINY-Gastric04 (NCT04704934)

The efficacy and safety of Enhertu were evaluated in DESTINY-Gastric04, a Phase 3, randomised, multicentre, open-label, active-controlled study. The study included adult patients with HER2-positive locally advanced, unresectable, or metastatic gastric or GEJ adenocarcinoma who had progressed on or after a trastuzumab-containing regimen. Patients were randomised 1:1 to receive either Enhertu (N=246) or ramucirumab plus paclitaxel (N=248). Randomisation was stratified by HER2 status (IHC 3+ or IHC 2+/ISH-positive), geographical region (Asia [excluding mainland China] versus Western Europe versus mainland China/rest of the world), and time to progression on first-line therapy (<6 months or ≥6 months). Tumour samples were required to have locally or centrally confirmed HER2 positivity defined as IHC 3+ or IHC 2+/ISH-positive. The study excluded patients with a history of ILD/pneumonitis requiring treatment with steroids or ILD/pneumonitis at screening, patients with a history of clinically significant cardiac disease, and patients with active brain metastases. Treatment was administered until disease progression, death, or unacceptable toxicity. The primary efficacy outcome measure was overall survival (OS). PFS, confirmed ORR, and DOR were secondary outcome measures.

Demographic and baseline disease characteristics were similar between treatment arms. Of the 494 patients enrolled in DESTINY-Gastric04, the median age was 63.7 years (range 21.1 to 87.0); 79.4% were male; 49.8% were White, 40.1% were Asian, and 0.4% were Black or African American. Patients had an ECOG performance status of either 0 (37.4%) or 1 (61.9%); 61.1% had gastric adenocarcinoma and 38.9% had GEJ adenocarcinoma; 84% were IHC 3+ and 16% were IHC 2+/ISH-positive; 70% of patients had two or more metastatic sites, 61.7% had liver metastases, 6.9% had brain metastases; 15.6% of patients previously received immunotherapy.

Efficacy results are summarised in Table 11 and Figure 10.

Table 11: Efficacy results in DESTINY-Gastric04

Table 11. Efficacy Results in DESTINY-Gastric04		
Efficacy Parameter	Enhertu N=246	Ramucirumab plus paclitaxel N=248
Overall Survival (OS)		
Number of events (%)	124 (50.4)	142 (57.3)
Median, months (95% CI)	14.7 (12.1, 16.6)	11.4 (9.9, 15.5)
Hazard ratio (95% CI)*	0.70 (0.55, 0.90)	
p-value†	P=0.0044	
Progression-free Survival (PFS) per Investigator Assessment		
Number of events (%)	166 (67.5)	156 (62.9)
Median, months (95% CI)	6.7 (5.6, 7.1)	5.6 (4.9, 5.8)
Hazard ratio (95% CI)*	0.74 (0.59, 0.92)	
p-value†	p=0.0074	
Confirmed Objective Response Rate (ORR) per Investigator Assessment††		
n (%)	104 (44.3)	69 (29.1)
95% CI	(37.8, 50.9)	(23.4, 35.3)
p-value§	p=0.0006	
Complete Response n (%)	7 (3.0)	3 (1.3)
Partial Response n (%)	97 (41.3)	66 (27.8)
Duration of Response (DOR) per Investigator Assessment		
Median, months (95% CI)	7.4 (5.7, 10.1)	5.3 (4.1, 5.7)

CI=confidence interval

*Two-sided p-value from stratified log-rank test and stratified Cox proportional hazards model adjusted for IRT stratification factors: HER2 status (IHC 3+ or IHC 2+/ISH+).


†Based on log rank test stratified by HER2 status (IHC3+ or IHC2+/ISH+)

††ORR-eligible subjects are those who were randomised at least 77 days (i.e., 2 × 6 weeks - 1 week) before DCO date of interim analysis. Confirmed ORR is calculated using the eligible subjects as the denominator:

Enhertu = 235, ramucirumab plus paclitaxel = 237

§p-value for the difference in ORR uses the Cochran-Mantel-Haenszel test adjusted for stratification factor: HER2 status (IHC 3+ or IHC 2+/ISH+).

Figure 10: Kaplan-Meier plot of overall survival (full analysis set)



DESTINY-Gastric02 (NCT04014075)

The efficacy and safety of Enhertu were studied in DESTINY-Gastric02, a Phase 2, multicentre, open-label, single-arm study conducted at sites in Europe and the United States. The study enrolled patients with locally advanced or metastatic HER2-positive gastric or GEJ adenocarcinoma who had progressed on a prior trastuzumab-based regimen. Patients were required to have centrally confirmed HER2 positivity defined as IHC 3+ or IHC 2+/ISH-positive. The study excluded patients with a history of ILD/pneumonitis requiring treatment with steroids or ILD/pneumonitis at screening, patients with a history of clinically significant cardiac disease, and patients with active brain metastases. Enhertu was administered by intravenous infusion at 6.4 mg/kg every three weeks until disease progression, death, withdrawal of consent, or unacceptable toxicity. The primary efficacy outcome measure was confirmed ORR assessed by ICR based on RECIST v1.1. DOR and OS were secondary endpoints.

Of the 79 patients enrolled in DESTINY-Gastric02, demographic and baseline disease characteristics were: median age 61 years (range 20 to 78); 72% were male; 87% were White, 5.0% were Asian and 1.0% were Black or African American. Patients had an ECOG performance status of either 0 (37%) or 1 (63%); 34% had gastric adenocarcinoma and 66% had GEJ adenocarcinoma; 86% were IHC 3+ and 13% were IHC 2+/ISH-positive, and 63% had liver metastases.

Efficacy results for ORR and DOR are summarised in Table 12.

Table 12: Efficacy results in DESTINY-Gastric02 (full analysis set*)

Efficacy parameter	DESTINY-Gastric02 N = 79
<i>Data cutoff date 08 November 2021</i>	
Confirmed objective response rate[†] % (95% CI) [‡]	41.8 (30.8, 53.4)

Efficacy parameter	DESTINY-Gastric02 N = 79
<i>Data cutoff date 08 November 2021</i>	
Complete response n (%)	4 (5.1)
Partial response n (%)	29 (36.7)
Duration of response Median [§] , months (95% CI) [¶]	8.1 (5.9, NE)

NE = Not estimable

*Includes all patients who received at least one dose of Enhertu

†Assessed by independent central review

‡Calculated using Clopper-Pearson method

§Based on Kaplan-Meier estimate

¶Calculated using the Brookmeyer and Crowley method

DESTINY-Gastric01 (NCT03329690)

The efficacy and safety of Enhertu were studied in DESTINY-Gastric01, a Phase 2, multicentre, open-label, randomised study conducted at sites in Japan and South Korea. This supportive study included adult patients with locally advanced or metastatic HER2-positive gastric or GEJ adenocarcinoma who had progressed on at least two prior regimens, including trastuzumab, a fluoropyrimidine agent, and a platinum agent. Patients were randomised 2:1 to receive either Enhertu (N = 126) or physician's choice of chemotherapy: either irinotecan (N = 55) or paclitaxel (N = 7). Tumour samples were required to have centrally confirmed HER2 positivity defined as IHC 3+ or IHC 2+/ISH-positive. The study excluded patients with a history of ILD/pneumonitis requiring treatment with steroids or ILD/pneumonitis at screening, patients with a history of clinically significant cardiac disease, and patients with active brain metastases. Treatment was administered until disease progression, death, withdrawal of consent, or unacceptable toxicity. The primary efficacy outcome measure was unconfirmed ORR assessed by ICR based on RECIST v1.1. Overall survival (OS), progression-free survival (PFS), DOR, and confirmed ORR were secondary outcome measures.

Demographic and baseline disease characteristics were similar between treatment arms. Of the 188 patients, the median age was 66 years (range 28 to 82); 76% were male; 100% were Asian. Patients had an ECOG performance status of either 0 (49%) or 1 (51%); 87% had gastric adenocarcinoma and 13% had GEJ adenocarcinoma; 76% were IHC 3+ and 23% were IHC 2+/ISH-positive; 54% had liver metastases; 29% had lung metastases; the sum of diameters of target lesions was < 5 cm in 47%, ≥ 5 to < 10 cm in 30%, and ≥ 10 cm in 17%; 55% had two and 45% had three or more prior regimens in the locally advanced or metastatic setting.

Efficacy results (data cutoff date: 03 June 2020) for Enhertu (n = 126) vs. physician's choice of chemotherapy (n = 62) were confirmed ORR 39.7% (95% CI: 31.1, 48.8) vs. 11.3% (95% CI: 4.7, 21.9). Complete response rate was 7.9% vs. 0% and partial response rate was 31.7% vs. 11.3%. Additional efficacy results for Enhertu vs. physician's choice of chemotherapy were median DOR of 12.5 months (95% CI: 5.6, NE) vs. 3.9 months (95% CI: 3.0, 4.9). Median PFS was 5.6 months (95% CI: 4.3, 6.9) vs. 3.5 months (95% CI: 2.0, 4.3; hazard ratio = 0.47 [95% CI: 0.31, 0.71]). An OS analysis, prespecified at 133 deaths, showed survival benefit with Enhertu treatment compared to the physician's choice group (hazard ratio = 0.60). The median OS was 12.5 months (95% CI: 10.3, 15.2) in the Enhertu group and 8.9 months (95% CI: 6.4, 10.4) in the physician's choice group.

Other Unresectable or Metastatic Solid Tumours

The efficacy and safety of Enhertu were evaluated in adult patients with previously treated unresectable or metastatic HER2-positive solid tumours who were enrolled in one of three Phase 2, multicentre studies: DESTINY-PanTumor02, and supportive studies DESTINY-Lung01, and DESTINY-CRC02.

DESTINY-PanTumor02

DESTINY-PanTumor02 was a Phase 2, multicentre, open-label study that enrolled and treated 267 adult patients with locally advanced, unresectable, or metastatic HER2-expressing (IHC3+, IHC2+) solid tumours who have progressed following at least one prior systemic regimen or who have no satisfactory alternative treatment option. Patients received Enhertu 5.4 mg/kg by intravenous infusion every three weeks. Treatment was administered until disease progression, death, withdrawal of consent, or unacceptable toxicity. The study excluded patients with a history of ILD/pneumonitis requiring treatment with steroids or ILD/pneumonitis at screening and clinically significant cardiac disease. Patients were also excluded for untreated and symptomatic brain metastases or ECOG performance status >1.

Efficacy was assessed *post-hoc* in the subgroup of patients (N = 111) with HER2-positive (IHC3+ by either local or central assessment) solid tumours. The primary efficacy outcome measure was investigator-assessed confirmed objective response rate (ORR) and a secondary efficacy outcome measure was duration of response (DOR). Independent central review (ICR) based on RECIST v1.1 was applied retrospectively in the study.

Of the 111 patients, demographic and baseline disease characteristics were: median age 64 years (range 23 to 85); 59.5% were female; 57.7% were White, 34.2% were Asian, and 3.6% were Black or African American. Patients had an ECOG performance status of either 0 (48.6%) or 1 (51.4%) at baseline. The median number of prior regimens was 2 (range: 0 to 9).

Efficacy results for HER2-positive (IHC3+) solid tumours are summarised in Table 13 and Table 14.

Table 13: Efficacy results in HER2-positive (IHC3+) patients in DESTINY-PanTumor02

Efficacy Parameter	DESTINY-PanTumor02 N = 111
Confirmed Objective Response Rate[†]	
n (%)	58 (52.3)
95% CI [‡]	(42.6, 61.8)
Complete Response, n (%)	12 (10.8)
Partial Response, n (%)	46 (41.4)
Duration of Response[†]	
Median [§] , months (95% CI) [¶]	21.1 (10.6, 25.0)

CI = Confidence interval

[†]Assessed by investigator

[‡]CI is derived based on the Clopper-Pearson method

[§]Calculated using the Kaplan-Meier technique

[¶]CI is derived based on the Brookmeyer-Crowley method.

Table 14: Efficacy results in HER2-positive (IHC3+) patients by tumour type in DESTINY-PanTumor02

Tumour Type	Patients N	Confirmed ORR [†] % (95% CI) [‡]	DOR Median [§] , months (95% CI) [‡]
Bladder cancer	27	44.4 (25.5, 64.7)	8.7 (4.1, 10.6)
Biliary tract cancer	22	40.9 (20.7, 63.6)	8.6 (2.1, NE)
Endometrial cancer	16	62.5 (35.4, 84.8)	28.8 (4.2, NE)
Ovarian cancer	15	73.3 (44.9, 92.2)	22.1 (2.8, NE)
Cervical cancer	10	80.0 (44.4, 97.5)	NE (9.3, NE)
Salivary gland cancer	9	55.6 (21.2, 86.3)	10.9 (4.1, NE)
Pancreatic cancer	5	0.0 (0.0, 52.2)	NE

Extramammary Paget's disease	1	100.0 (2.5, 100.0)	23.6 (NE, NE)
Head and neck	1	100.0 (2.5, 100.0)	NE
Lip and/or oral cavity cancer	1	0.0 (0, 97.5)	NE
Oesophageal adenocarcinoma	1	100 (2.5, 100)	NE
Oesophageal squamous cell carcinoma	1	0.0 (0, 97.5)	NE
Oropharyngeal neoplasm	1	100.0 (2.5, 100.0)	21.1 (NE, NE)
Vulvar cancer	1	100.0 (2.5, 100.0)	NE

CI = Confidence interval, NE = Not estimable

†Assessed by investigator

*CI is derived based on the Clopper-Pearson method

§Calculated using the Kaplan-Meier technique

Supportive Studies

DESTINY-Lung01 was a Phase 2, multicentre, open-label, 2-cohort study that enrolled and treated 181 patients with previously treated, unresectable or metastatic, HER2-expressing (IHC3+ or IHC2+) or HER2-mutant non-small cell lung cancer (NSCLC). Patients received Enhertu 5.4 mg/kg by intravenous infusion every three weeks. Efficacy was assessed in the subgroup of patients (N = 17) with centrally confirmed HER2-positive (IHC3+) non-small cell lung cancer. The primary efficacy outcome of confirmed ORR was 52.9% (95% CI 27.8, 77.0) and the key secondary efficacy outcome measure of DOR was 6.9 months (95% CI 4.0, 9.8). All outcomes were assessed by independent central review (ICR) based on RECIST v1.1.

DESTINY-CRC02 was a Phase 2, multicentre, randomised, 2-arm study that enrolled and treated 122 patients with previously treated, unresectable or metastatic HER2-expressing (IHC3+ or IHC2+/ISH+) colorectal cancer (CRC). Patients received Enhertu 5.4 mg/kg by intravenous infusion every three weeks. Efficacy was assessed in the subgroup of patients (N = 64) with centrally confirmed HER2-positive (IHC3+) CRC. The primary efficacy outcome measure of confirmed ORR was 46.9% (95% CI 34.3, 59.8) and the key secondary efficacy outcome measure of DOR was 5.5 months (95% CI 4.2, 8.1). All outcomes were assessed by independent central review (ICR) based on RECIST v1.1.

Paediatric population

The European Medicines Agency has waived the obligation to submit the results of studies in all subsets of the paediatric population in breast cancer, NSCLC and gastric cancer (see section 4.2 for information on paediatric use).

This medicinal product has been authorised under a so-called 'conditional approval' scheme. This means that further evidence on this medicinal product is awaited.

The European Medicines Agency will review new information on this medicinal product at least every year and this SmPC will be updated as necessary.

5.2 Pharmacokinetic properties

At the recommended dosage of Enhertu for patients with metastatic breast cancer, NSCLC, and HER2-positive (IHC 3+) solid tumours (5.4 mg/kg), the geometric mean (coefficient of variation [CV]%) C_{max} at steady state of trastuzumab deruxtecan and topoisomerase I inhibitor, DXd, were 131 µg/mL (22%) and 5.1 ng/mL (47%), respectively, and the AUC at steady state of trastuzumab deruxtecan and DXd were 739 µg day/mL (34%) and 30.7 ng day/mL (49%), respectively.

At the recommended dosage of Enhertu for patients with HER2-positive gastric cancer (6.4 mg/kg), the geometric mean C_{max} at steady state of trastuzumab deruxtecan and DXd were 131 µg/mL (21%)

and 5.2 ng/mL (42.0%), respectively, and the AUC at steady state of trastuzumab deruxtecan and DXd were 762 µg day/mL (29%) and 32.1 ng day/mL (44.1%), respectively.

Absorption

Trastuzumab deruxtecan is administered intravenously. There have been no studies performed with other routes of administration.

Distribution

Based on population pharmacokinetic analysis, the volume of distribution of the central compartment (V_c) of trastuzumab deruxtecan and DXd, were estimated to be 2.8 L and 28.0 L, respectively. The volume of distribution of the peripheral compartment (V_p) of trastuzumab deruxtecan was estimated to be 6.0 L.

In vitro, the mean human plasma protein binding of DXd was approximately 97%.

In vitro, the blood to plasma concentration ratio of DXd was approximately 0.6.

Biotransformation

Trastuzumab deruxtecan undergoes intracellular cleavage by lysosomal enzymes to release the DXd.

The humanised HER2 IgG1 monoclonal antibody is expected to be degraded into small peptides and amino acids via catabolic pathways in the same manner as endogenous IgG.

In vitro metabolism studies in human liver microsomes indicate that DXd is metabolised mainly by CYP3A4 via oxidative pathways and does not undergo significant metabolism by uridine diphosphate-glucuronosyltransferases (UGT) or other CYP enzymes.

Elimination

Following intravenous administration of trastuzumab deruxtecan in patients with metastatic HER2-positive, HER2-low breast cancer or HER2-mutant NSCLC, the clearance of trastuzumab deruxtecan in population pharmacokinetic analysis was calculated to be 0.4 L/day and the clearance of DXd was 18.4 L/h. In patients with locally advanced or metastatic gastric or GEJ adenocarcinoma, trastuzumab deruxtecan clearance was approximately 20% higher than in patients with metastatic HER2-positive breast cancer. In cycle 3, the apparent elimination half-life (t_{1/2}) of trastuzumab deruxtecan and released DXd was approximately 7 days. Moderate accumulation (approximately 35% in cycle 3 compared to cycle 1) of trastuzumab deruxtecan was observed.

Following intravenous administration of DXd to rats, the major excretion pathway was faeces via the biliary route. DXd was the most abundant component in urine, faeces, and bile. Following single intravenous administration of trastuzumab deruxtecan (6.4 mg/kg) to monkeys, unchanged released DXd was the most abundant component in urine and faeces. DXd excretion was not studied in humans.

In vitro interactions

Effects of Enhertu on the pharmacokinetics of other medicinal products

In vitro studies indicate DXd does not inhibit major CYP450 enzymes including CYP1A2, 2B6, 2C8, 2C9, 2C19, 2D6 and 3A. *In vitro* studies indicate that DXd does not inhibit OAT1, OAT3, OCT1, OCT2, OATP1B1, OATP1B3, MATE1, MATE2-K, P-gp, BCRP, or BSEP transporters.

Effects of other medicinal products on the pharmacokinetics of Enhertu

In vitro, DXd was a substrate of P-gp, OATP1B1, OATP1B3, MATE2-K, MRP1, and BCRP.

No clinically meaningful interaction is expected with medicinal products that are inhibitors of MATE2-K, MRP1, P-gp, OATP1B, or BCRP transporters (see section 4.5).

Linearity/non-linearity

The exposure of trastuzumab deruxtecan and released DXd when administered intravenously increased in proportion to dose in the 3.2 mg/kg to 8.0 mg/kg dose range (approximately 0.6 to 1.5 times the recommended dose) with low to moderate inter-subject variability. Based on population pharmacokinetic analysis, inter-subject variability in trastuzumab deruxtecan and DXd elimination clearances were 24% and 28%, respectively, and for central volume of distribution were 16% and 55%, respectively. The intra-subject variability in trastuzumab deruxtecan and DXd AUC values (area under the serum concentration versus time curve) was approximately 8% and 14%, respectively.

Special populations

Based on population pharmacokinetic analysis, age (20-96 years), race, ethnicity, sex and body weight did not have a clinically meaningful effect on exposure of trastuzumab deruxtecan or released DXd.

Elderly

The population PK analysis showed that age (range: 20-96 years) did not affect the PK of trastuzumab deruxtecan.

Renal impairment

No dedicated renal impairment study was conducted. Based on population pharmacokinetic analysis including patients with mild (creatinine clearance [CL_{Cr}] ≥ 60 and < 90 mL/min) or moderate (CL_{Cr} ≥ 30 and < 60 mL/min) renal impairment (estimated by Cockcroft-Gault), the pharmacokinetics of the released DXd was not affected by mild or moderate renal impairment as compared to normal renal function (CL_{Cr} ≥ 90 mL/min).

Hepatic impairment

No dedicated hepatic impairment study was conducted. Based on population pharmacokinetic analysis, the impact of changes on pharmacokinetics of trastuzumab deruxtecan in patients with total bilirubin ≤ 1.5 times ULN, irrespective of AST level, is not clinically meaningful. There is limited data for patients with total bilirubin > 1.5 to 3 times ULN, irrespective of AST level, to draw conclusions, and no data is available for patients with total bilirubin > 3 times ULN, irrespective of AST level (see sections 4.2 and 4.4).

Paediatric population

No studies have been conducted to investigate the pharmacokinetics of trastuzumab deruxtecan in children or adolescents.

5.3 Preclinical safety data

In animals, toxicities were observed in lymphatic and haematopoietic organs, intestines, kidneys, lungs, testes and skin following the administration of trastuzumab deruxtecan at exposure levels of the topoisomerase I inhibitor (DXd) below clinical plasma exposure. In these animals, antibody-drug conjugate (ADC) exposure levels were similar or above clinical plasma exposure.

DXd was clastogenic in both an *in vivo* rat bone marrow micronucleus assay and an *in vitro* Chinese hamster lung chromosome aberration assay and was not mutagenic in an *in vitro* bacterial reverse mutation assay.

Carcinogenicity studies have not been conducted with trastuzumab deruxtecan.

Dedicated fertility studies have not been conducted with trastuzumab deruxtecan. Based on results from general animal toxicity studies, trastuzumab deruxtecan may impair male reproductive function and fertility.

There were no animal reproductive or developmental toxicity studies conducted with trastuzumab deruxtecan. Based on results from general animal toxicity studies, trastuzumab deruxtecan and DXd were toxic to rapidly dividing cells (lymphatic/haematopoietic organs, intestine, or testes), and DXd was genotoxic, suggesting the potential for embryotoxicity and teratogenicity.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

L-histidine
L-histidine hydrochloride monohydrate
Sucrose
Polysorbate 80 (E433)

6.2 Incompatibilities

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products except those mentioned in section 6.6.

Sodium chloride solution for infusion must not be used for reconstitution or dilution since it may cause particulate formation.

6.3 Shelf life

Unopened vial

4 years.

Reconstituted solution

Chemical and physical in-use stability has been demonstrated for up to 48 hours at 2 °C to 8 °C.

From a microbiological point of view, the product should be used immediately. If not used immediately, in-use storage times and conditions prior to use are the responsibility of the user and would normally not be longer than 24 hours at 2 °C to 8 °C, unless reconstitution has taken place in controlled and validated aseptic conditions.

Diluted solution

It is recommended that the diluted solution be used immediately. If not used immediately, the reconstituted solution diluted in infusion bags containing 5% glucose solution may be stored at room temperature (≤ 30 °C) for up to 4 hours including preparation and infusion or in a refrigerator at 2 °C to 8 °C for up to 24 hours, protected from light.

6.4 Special precautions for storage

Store in a refrigerator (2 °C - 8 °C).

Do not freeze.

Store in the original carton in order to protect from light.

For storage conditions after reconstitution and dilution of the medicinal product, see section 6.3.

6.5 Nature and contents of container

Enhertu is provided in 10 mL Type 1 amber borosilicate glass vial sealed with a fluoro-resin laminated butyl rubber stopper, and a polypropylene/aluminium yellow flip-off crimp cap. Each carton contains 1 vial.

6.6 Special precautions for disposal and other handling

In order to prevent medicinal product errors, it is important to check the vial labels to ensure that the medicinal product being prepared and administered is Enhertu (trastuzumab deruxtecan) and not trastuzumab or trastuzumab emtansine.

Appropriate procedures for the preparation of chemotherapeutic medicinal products should be used. Appropriate aseptic technique should be used for the following reconstitution and dilution procedures.

Reconstitution

- Reconstitute immediately before dilution.
- More than one vial may be needed for a full dose. Calculate the dose (mg), the total volume of reconstituted Enhertu solution required, and the number of vial(s) of Enhertu needed (see section 4.2).
- Reconstitute each 100 mg vial using a sterile syringe to slowly inject 5 mL of water for injection into each vial to obtain a final concentration of 20 mg/mL.
- Swirl the vial gently until completely dissolved. Do not shake.
- From a microbiological point of view, the product should be used immediately. If not used immediately, chemical and physical in-use stability has been demonstrated for up to 48 hours at 2 °C to 8 °C. Store the reconstituted Enhertu vials in a refrigerator at 2 °C to 8 °C, protected from light. Do not freeze.
- The reconstituted product contains no preservative and is intended for single use only.

Dilution

- Withdraw the calculated amount from the vial(s) using a sterile syringe. Inspect the reconstituted solution for particulates and discolouration. The solution should be clear and colourless to light yellow. Do not use if visible particles are observed or if the solution is cloudy or discoloured.
- Dilute the calculated volume of reconstituted Enhertu in an infusion bag containing 100 mL of 5% glucose solution for infusion. Do not use sodium chloride solution (see section 6.2). An infusion bag made of polyvinylchloride or polyolefin (copolymer of ethylene and polypropylene) is recommended.
- Gently invert the infusion bag to thoroughly mix the solution. Do not shake.
- Cover the infusion bag to protect from light.
- If not used immediately, store at room temperature (≤ 30 °C) for up to 4 hours including preparation and infusion or in a refrigerator at 2 °C to 8 °C for up to 24 hours, protected from light. Do not freeze.
- Discard any unused portion left in the vial.

Administration

- If the prepared infusion solution was stored refrigerated (2 °C to 8 °C), it is recommended that the solution be allowed to equilibrate to room temperature prior to administration, protected from light.
- Administer Enhertu as an intravenous infusion only with a 0.20 or 0.22 micron in-line polyethersulfone (PES) or polysulfone (PS) filter.

- The initial dose should be administered as a 90-minute intravenous infusion. If the prior infusion was well tolerated, subsequent doses of Enhertu may be administered as 30-minute infusions. Do not administer as an intravenous push or bolus (see section 4.2).
- Cover the infusion bag to protect from light.
- Do not mix Enhertu with other medicinal products or administer other medicinal products through the same intravenous line.

Disposal

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Daiichi Sankyo Europe GmbH
Zielstattstrasse 48
81379 Munich
Germany

8. MARKETING AUTHORISATION NUMBER

EU/1/20/1508/001

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 18 January 2021
Date of latest renewal: 21 November 2025

10. DATE OF REVISION OF THE TEXT

28 August 2026

Detailed information on this medicinal product is available on the website of the European Medicines Agency <https://www.ema.europa.eu>