

1. NAME OF THE MEDICINAL PRODUCT

Fludara 10 mg film-coated tablets

2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains 10mg fludarabine phosphate.

Excipients: Lactose monohydrate 74.75 mg

For the full list of excipients, see section 6.1

3. PHARMACEUTICAL FORM

Film-coated tablets.

Salmon-pink, capsule-shaped tablet marked with 'LN' in a regular hexagon on one side.

4. CLINICAL PARTICULARS

4.1 Therapeutic indications

Treatment of B-cell chronic lymphocytic leukaemia (CLL) in adult patients with sufficient bone marrow reserves.

First line treatment with Fludara should only be initiated in adult patients with advanced disease, Rai stages III/IV (Binet stage C) or Rai stages I/II (Binet stage A/B) where the patient has disease related symptoms or evidence of progressive disease.

4.2 Posology and method of administration

Posology

The recommended dose is 40 mg fludarabine phosphate/m² body surface given daily for 5 consecutive days every 28 days by oral route. This dose corresponds to 1.6 times the recommended intravenous dose of fludarabine phosphate (25 mg/m² body surface per day).

The following table provides guidance for determining the number of tablets of Fludara to be administered:

Body Surface Area (BSA) [m ²]	Calculated total daily dose based on BSA (rounded up or down to whole number) [mg/day]	Number of tablets per day (total daily dose)
0.75 - 0.88	30 – 35	3 (30 mg)
0.89 - 1.13	36 – 45	4 (40 mg)
1.14 - 1.38	46 – 55	5 (50 mg)
1.39 - 1.63	56 – 65	6 (60 mg)
1.64 - 1.88	66 – 75	7 (70 mg)
1.89 - 2.13	76 – 85	8 (80 mg)
2.14 - 2.38	86 – 95	9 (90 mg)
2.39 - 2.50	96 – 100	10 (100 mg)

The duration of treatment depends on the success of treatment and the tolerability of the drug. Fludara should be administered until best response is achieved (complete or partial remission, usually 6 cycles) and then the drug should be discontinued.

Dose adjustments for the first treatment cycle (start of therapy with Fludara) are not recommended (except in patients with impairment of renal function, see 'Patients with renal impairment').

Patients undergoing treatment with Fludara should be closely monitored for response and toxicity.

Individual dosing should be carefully adjusted according to the observed haematological toxicity.

If at the start of a subsequent cycle cell numbers are too low to administer the recommended dosage and there is evidence of treatment associated myelosuppression, the planned treatment cycle should be postponed until granulocyte count is above $1.0 \times 10^9/\text{L}$ and platelet count is above $100 \times 10^9/\text{L}$. Treatment should only be postponed up to a maximum of two weeks. If granulocyte and platelet counts have not recovered after two weeks of postponement, the dose should be reduced according to the suggested dose adjustments in the table below.

Granulocytes and / or [$10^9/\text{L}$]		Platelets	Fludarabine phosphate dose
0.5 - 1.0		50 - 100	30 mg/m ² /day
<0.5		<50	20 mg/m ² /day

Dose should not be reduced if thrombocytopenia is disease related.

If a patient does not respond to treatment after two cycles and shows no or little haematological toxicity a careful dose adjustment towards higher fludarabine phosphate doses in subsequent treatment cycles could be considered.

Patients with renal impairment

Doses should be adjusted for patients with reduced kidney function. If creatinine clearance is between 30 and 70 ml/min, the dose should be reduced by up to 50% and close haematological monitoring should be used to assess toxicity (see section 4.4).

Fludara treatment is contraindicated if creatinine clearance is < 30 ml/min (see section 4.3).

Patients with hepatic impairment

No data are available concerning the use of Fludara in patients with hepatic impairment. In this group of patients, Fludara should be used with caution.

Paediatric population

The safety and efficacy of Fludara in children below the age of 18 years have not been established. Therefore, Fludara is not recommended for use in children.

Older people

Since there are limited data for the use of Fludara in older people (> 75 years), caution should be exercised with the administration of Fludara in these patients.

In patients over the age of 65 years, creatinine clearance should be measured (see "Patients with renal impairment" and section 4.4).

Method of administration

Fludara should be prescribed by a qualified physician experienced in the use of antineoplastic therapy.

Fludara can be taken either on an empty stomach or together with food. The tablets have to be swallowed whole with water, they should not be chewed or broken.

Precautions to be taken before handling the medicinal product

For instructions on handling of the medicinal product, see section 6.6.

4.3 Contraindications

- Hypersensitivity to the active substance or to any of the excipients listed in section 6.1.
- Renal impairment with creatinine clearance <30 ml/min.
- Decompensated haemolytic anaemia.
- Lactation.

4.4 Special warnings and precautions for use

Myelosuppression

Severe bone marrow suppression, notably anaemia, thrombocytopenia and neutropenia, has been reported in patients treated with Fludara. In a Phase I intravenous study in adult solid tumour patients, the median time to nadir counts was 13 days (range 3 – 25 days) for granulocytes and 16 days (range 2 - 32 days) for platelets. Most patients had haematologic impairment at baseline either as a result of disease or as a result of prior myelosuppressive therapy.

Cumulative myelosuppression may be seen. While chemotherapy-induced myelosuppression is often reversible, administration of fludarabine phosphate requires careful haematologic monitoring.

Fludarabine phosphate is a potent antineoplastic agent with potentially significant toxic side effects. Patients undergoing therapy should be closely observed for signs of haematologic and non-haematologic toxicity. Periodic assessment of peripheral blood counts is recommended to detect the development of anaemia, neutropenia and thrombocytopenia.

Several instances of trilineage bone marrow hypoplasia or aplasia resulting in pancytopenia, sometimes resulting in death, have been reported in adult patients. The duration of clinically significant cytopenia in the reported cases has ranged from approximately 2 months to approximately 1 year. These episodes have occurred both in previously treated or untreated patients.

As with other cytotoxics, caution should be exercised with fludarabine phosphate, when further haematopoietic stem cell sampling is considered.

Autoimmune disorders

Irrespective of any previous history of autoimmune processes or Coombs test status, life-threatening and sometimes fatal autoimmune phenomena (see section 4.8) have been reported to occur during or after treatment with Fludara. The majority of patients experiencing haemolytic anaemia developed a recurrence in the haemolytic process after rechallenge with Fludara.

Patients treated with Fludara should be closely monitored for signs of haemolysis.

Discontinuation of therapy with Fludara is recommended in case of haemolysis. Blood transfusion (irradiated, see below) and adrenocorticoid preparations are the most common treatment measures for autoimmune haemolytic anaemia.

Neurotoxicity

The effect of chronic administration of Fludara on the central nervous system is unknown. However, patients tolerated the recommended intravenous dose, in some studies for relatively long treatment times (for up to 26 courses of therapy). Patients should be closely observed for signs of neurologic effects.

When used at high doses in dose-ranging studies in patients with acute leukaemia, intravenous Fludara was associated with severe neurological effects, including blindness, coma and death. Symptoms appeared from 21 to 60 days from last dose. This severe central nervous system toxicity occurred in 36 % of patients treated intravenously with doses approximately four times greater (96 mg/m²/day for 5 - 7 days) than the recommended dose. In patients treated at doses in the range of the dose recommended for CLL, severe central nervous system toxicity occurred rarely (coma, seizures and agitation) or uncommonly (confusion) (see section 4.8).

In post-marketing experience neurotoxicity has been reported to occur earlier or later than in clinical trials.

Administration of Fludara can be associated with leukoencephalopathy (LE), acute toxic leukoencephalopathy (ATL) or reversible posterior leukoencephalopathy syndrome (RPLS). These may occur:

- at the recommended dose
 - when Fludara is given following, or in combination with, medications known to be associated with LE, ATL or RPLS,
 - or when Fludara is given in patients with other risk factors such as cranial or total body irradiation, Hematopoietic Cell Transplantation, Graft versus Host Disease, renal impairment, or hepatic encephalopathy.
- at doses higher than the recommended dose

LE, ATL or RPLS symptoms may include headache, nausea and vomiting, seizures, visual disturbances such as vision loss, altered sensorium, and focal neurological deficits. Additional effects may include optic neuritis, and papillitis, confusion, somnolence, agitation, paraparesis/ quadriparesis, muscle spasticity and incontinence.

LE/ ATL/ RPLS may be irreversible, life-threatening, or fatal.

Whenever LE, ATL or RPLS is suspected, fludarabine treatment should be stopped. Patients should be monitored and should undergo brain imaging, preferably utilizing MRI. If the diagnosis is confirmed, fludarabine therapy should be permanently discontinued.

Tumour lysis syndrome

Tumour lysis syndrome has been reported in CLL patients with large tumour burdens. Since Fludara can induce a response as early as the first week of treatment, precautions should be taken in those patients at risk of developing this complication, and hospitalisation may be recommended for these patients during the first course of treatment.

Transfusion-associated graft-versus-host disease

Transfusion-associated graft-versus-host disease (reaction by the transfused immunocompetent lymphocytes to the host) has been observed after transfusion of non-irradiated blood in Fludara-treated patients. Fatal outcome as a consequence of this disease has been reported with a high frequency. Therefore, to minimise the risk of transfusion-associated graft-versus-host disease, patients who require blood transfusion and who are undergoing, or who have received treatment with Fludara should receive irradiated blood only.

Skin cancer

The worsening or flare up of pre-existing skin cancer lesions as well as new onset of skin cancer has been reported in some patients during or after Fludara therapy.

Impaired state of health

In patients with impaired state of health, Fludara should be given with caution and after careful risk/benefit consideration. This applies especially for patients with severe impairment of bone marrow function (thrombocytopenia, anaemia and/or granulocytopenia), immunodeficiency or with a history of opportunistic infection.

Renal impairment

The total body clearance of the principle plasma metabolite 2-F-ara-A shows a correlation with creatinine clearance, indicating the importance of the renal excretion pathway for the elimination of the compound. Patients with reduced renal function demonstrated an increased total body exposure (AUC of 2F-ara-A). There are limited clinical data available in patients with impairment of renal function (creatinine clearance < 70 ml/min).

Fludara must be administered cautiously in patients with renal insufficiency. In patients with moderate impairment of renal function (creatinine clearance between 30 and 70 ml/min), the dose should be reduced by up to 50% and the patient should be monitored closely (see section 4.2). Fludara treatment is contraindicated if creatinine clearance is < 30ml/min (see section 4.3).

Older people

Since there are limited data for the use of Fludara in older people (> 75 years), caution should be exercised with the administration of Fludara in these patients.

In patients aged 65 years or older, creatinine clearance should be measured before start of treatment, see “Renal impairment” and section 4.2.

Pregnancy

Fludarabine phosphate has been shown to be genotoxic. Fludarabine phosphate has also been shown to be both embryotoxic and fetotoxic in rabbits and rats (see sections 5.3) Fludara may cause foetal harm when administered to pregnant females. Therefore, Fludara must not be used during pregnancy unless the potential benefit for the mother outweighs the potential risks to the foetus.

Females of childbearing potential receiving Fludara should be advised to avoid becoming pregnant, and to inform the treating physician immediately should this occur. (see sections 4.6 and 5.3)

Contraception in males and females

Due to the genotoxic risk of fludarabine phosphate, females of child-bearing potential must take effective contraceptive measures during and at least for 6 months after cessation of therapy. Male patients must use effective methods of contraception and be advised to not father a child while receiving Fludara, and at least for 3 months following completion of treatment (see section 4.6).

Vaccination

During and after treatment with Fludara vaccination with live vaccines should be avoided.

Retreatment options after initial Fludara treatment

A crossover from initial treatment with Fludara to chlorambucil for non-responders to Fludara should be avoided because most patients who have been resistant to Fludara have shown resistance to chlorambucil.

Change to Fludara IV

The reported incidence of nausea/vomiting was higher with the oral than the i.v. formulation. If this presents a persistent clinical problem it is recommended to switch to the i.v. formulation.

Excipients

Each Fludara 10 mg film-coated tablet contains 74.75 mg lactose monohydrate. Patients with rare hereditary problems of galactose intolerance, total lactase deficiency or glucose-galactose malabsorption should not take this medicine.

Fludara 10 mg film-coated tablet contains sodium. This medicine contains less than 1 mmol sodium (23 mg) per tablet, that is to say essentially 'sodium-free'.

4.5 Interaction with other medicinal products and other forms of interaction

In a clinical investigation using intravenous Fludara in combination with pentostatin (deoxycoformycin) for the treatment of refractory chronic lymphocytic leukaemia (CLL), there was an unacceptably high incidence of fatal pulmonary toxicity. Therefore, the use of Fludara in combination with pentostatin is not recommended.

Dipyridamole and other inhibitors of adenosine uptake may reduce the therapeutic efficacy of Fludara.

Clinical studies and in vitro experiments showed that during use of Fludara in combination with cytarabine the intracellular peak concentration and intracellular exposure of Ara-CTP (active metabolite of cytarabine) increased in leukaemic cells. Plasma concentrations of Ara-C and the elimination rate of Ara-CTP were not affected.

In a clinical investigation, pharmacokinetic parameters after peroral administration were not significantly affected by concomitant food intake (see section 5.2).

4.6 Fertility, pregnancy and lactation

Women of childbearing potential/Contraception in males and females.

Women of childbearing potential must be apprised of the potential hazard to the foetus.

Due to the genotoxic risk of fludarabine phosphate, women of childbearing potential must take effective contraceptive measures during and at least for 6 months after cessation of therapy. Male patients must use effective methods of contraception and be advised to not father a child while receiving Fludara, and at least for 3 months following completion of treatment.

Pregnancy

There are limited data from the use of fludarabine phosphate in pregnant women. Fludarabine phosphate has been shown to be genotoxic. Studies in animals have shown reproductive toxicity (see section 5.3). Fludara may cause foetal harm when taken by pregnant females. Therefore, Fludara must not be used during pregnancy, unless the potential benefit for the mother outweighs the potential risks to the foetus.

Women of childbearing potential receiving Fludara should be advised to avoid becoming pregnant, and to inform the treating physician immediately should this occur (see section 5.3).

Breast-feeding

It is not known whether fludarabine phosphates or its metabolites are excreted in human milk.

However, there is evidence from preclinical data that fludarabine phosphate and/or metabolites transfer from maternal blood to milk.

Because of the potential for serious adverse reactions to Fludara in breast-fed infants, Fludara is contraindicated during breast-feeding (see section 4.3).

Fertility

Fludara affects fertility in both males and females. Before Fludara treatment, patients planning pregnancy are advised to seek genetic counselling. Prior to Fludara treatment, male patients must seek advice on fertility preservation options.

4.7 Effects on ability to drive and use machines

Fludara may reduce the ability to drive and use machines, since e.g. fatigue, weakness, visual disturbances, confusion, agitation and seizures have been observed.

4.8 Undesirable effects

Summary of safety profile

Based on the experience with the use of Fludara, the most common adverse events include myelosuppression (neutropenia, thrombocytopenia and anaemia), infection including pneumonia, cough, fever, fatigue, weakness, nausea, vomiting and diarrhoea. Other commonly reported events include chills, oedema, malaise, peripheral neuropathy, visual disturbance, anorexia, mucositis, stomatitis and skin rash. Serious opportunistic infections have occurred in patients treated with Fludara. Fatalities as a consequence of serious adverse events have been reported.

Tabulated list of adverse reactions

The table below reports adverse events by MedDRA system organ classes (MedDRA SOCs). The frequencies are based on clinical trial data regardless of the causal relationship with Fludara. The rare adverse reactions were mainly identified from the post-marketing experience.

System Organ Class	Very Common (≥1/10)	Common (≥1/100 to <1/10)	Uncommon (≥1/1,000 to <1/100)	Rare (≥1/10,000 to <1/1,000)
Infections and infestations	Infections / Opportunistic infections (like latent viral reactivation, e.g. progressive multifocal leukoencephalopathy, Herpes zoster virus Epstein-Barr-virus), pneumonia			Lympho-proliferative disorder (EBV-associated)
Neoplasms benign, malignant and unspecified (incl cysts and polyps)		Myelodysplastic syndrome and Acute Myeloid Leukaemia (mainly associated with prior, concomitant or subsequent treatment with alkylating agents,		

		topoisomerase inhibitors or irradiation)		
Blood and lymphatic system disorders	Neutropenia, anaemia, thrombocytopenia	Myelosuppression		
Immune system disorders			Autoimmune disorder (including autoimmune haemolytic anaemia, Evan's syndrome, thrombocytopenic purpura, acquired haemophilia, pemphigus	
Metabolism and nutrition disorders		Anorexia	Tumour lysis syndrome (including renal failure, metabolic acidosis, hyperkalaemia, hypocalcaemia, hyperuricaemia, haematuria, urate crystalluria, hyperphosphataemia)	
Nervous system disorders		Peripheral neuropathy	Confusion	Coma, seizures, agitation
Eye disorders		Visual disturbances		Blindness, optic neuritis, optic neuropathy
Cardiac disorders				Heart failure, arrhythmia
Respiratory, thoracic and mediastinal disorders	Cough		Pulmonary toxicity (including pulmonary fibrosis, pneumonitis, dyspnoea)	
Gastro-intestinal disorders	Vomiting, diarrhoea, nausea	Stomatitis	Gastrointestinal haemorrhage, pancreatic enzymes abnormal	
Hepatobiliary			Hepatic	

disorders			enzymes abnormal	
Skin and subcutaneous tissue disorders		Rash		Skin cancer, necrolysis epidermal toxic (Lyell type) Stevens-Johnson syndrome
General disorders and administratio n site conditions	Fever, fatigue, weakness	Oedema, mucositis, chills, malaise		

The most appropriate MedDRA term to describe a certain adverse event is listed. Synonyms or related conditions are not listed, but should be taken into account as well. Adverse event term representation is based on MedDRA version 12.0.

Within each frequency grouping, undesirable effects are presented in order of decreasing seriousness.

Postmarketing experience with frequency unknown

- Nervous system disorders
 - Cerebral haemorrhage
 - Leukoencephalopathy (see section 4.4)
 - Acute toxic leukoencephalopathy (see section 4.4)
 - Reversible posterior leukoencephalopathy syndrome (RPLS) (see section 4.4)
- Respiratory, thoracic and mediastinal disorders
 - Pulmonary haemorrhage
- Renal and urinary disorder
 - Haemorrhagic cystitis

Reporting of suspected adverse reactions

Reporting suspected adverse reactions after authorisation of the medicinal product is important. It allows continued monitoring of the benefit/risk balance of the medicinal product. Healthcare professionals are asked to report any suspected adverse reactions via HPRA Pharmacovigilance, website www.hpra.ie.

4.9 Overdose

High doses of Fludara given intravenously have been associated with leukoencephalopathy, acute toxic leukoencephalopathy, or reversible posterior leukoencephalopathy syndrome (RPLS). Symptoms may include headache, nausea and vomiting, seizures, visual disturbances such as vision loss, altered sensorium, and focal neurological deficits. Additional effects may include optic neuritis, and papillitis, confusion, somnolence, agitation, paraparesis/ quadriparesis, muscle spasticity, incontinence, irreversible central nervous system toxicity characterised by delayed blindness, coma, and death. High doses are also associated with severe thrombocytopenia and neutropenia due to bone marrow suppression.

There is no known specific antidote for Fludara overdosage. Treatment consists of drug discontinuation and supportive therapy.

5. PHARMACOLOGICAL PROPERTIES

5.1 Pharmacodynamic properties

Pharmacotherapeutic group: Antineoplastic agents, purine analogues
ATC code: L01B B05

Mechanism of action

Fludara contains fludarabine phosphate, a water-soluble fluorinated nucleotide analogue of the antiviral agent vidarabine 9- β -D-arabinofuranosyladenine (ara-A) that is relatively resistant to deamination by adenosine deaminase.

Fludarabine phosphate is rapidly dephosphorylated to 2F-ara-A which is taken up by cells and then phosphorylated intracellularly by deoxycytidine kinase to the active triphosphate, 2F-ara-ATP. This metabolite has been shown to inhibit ribonucleotide reductase, DNA polymerase α/δ and ϵ , DNA primase and DNA ligase thereby inhibiting DNA synthesis. Furthermore, partial inhibition of RNA polymerase II and consequent reduction in protein synthesis occur.

While some aspects of the mechanism of action of 2F-ara-ATP are as yet unclear, it is assumed that effects on DNA, RNA and protein synthesis all contribute to inhibition of cell growth with inhibition of DNA synthesis being the dominant factor. In addition, in vitro studies have shown that exposure of CLL lymphocytes to 2F-ara-A triggers extensive DNA fragmentation and cell death characteristic of apoptosis.

Clinical efficacy and safety

A phase III trial in patients with previously untreated B-chronic lymphocytic leukaemia comparing treatment with Fludara vs. chlorambucil (40mg/m² q4 weeks) in 195 and 199 patients respectively showed the following outcome: statistically significant higher overall response rates and complete response rates after 1st line treatment with Fludara compared to chlorambucil (61.1% vs. 37.6% and 14.9% vs. 3.4%, respectively); statistically significant longer duration of response (19 vs. 12.2 months) and time to progression (17 vs. 13.2 months) for the patients in the Fludara group. The median survival of the two patient groups was 56.1 months for Fludara and 55.1 months for chlorambucil, a non-significant difference was also shown with performance status. The proportion of patients reported to have toxicities were comparable between Fludara patients (89.7%) and chlorambucil patients (89.9%). While the difference in the overall incidence of haematological toxicities was not significant between the two treatment groups, significantly greater proportions of Fludara patients experienced white blood cell (p=0.0054) and lymphocyte (p=0.0240) toxicities than chlorambucil patients. The proportions of patients who experienced nausea, vomiting and diarrhoea were significantly lower for Fludara patients (p<0.0001, p<0.0001, and p=0.0489, respectively) than chlorambucil patients. Toxicities of the liver were also reported for significantly (p=0.0487) less proportions of patients in the Fludara group than in the chlorambucil group. Patients who initially respond to Fludara have a chance of responding again to Fludara monotherapy.

A randomised trial of Fludara vs. cyclophosphamide, adriamycin and prednisone (CAP) in 208 patients with CLL Binet stage B or C revealed the following results in the subgroup of 103 previously treated patients: the overall response rate and the complete response rate were higher with Fludara compared to CAP (45% vs. 26% and 13% vs. 6%, respectively); response duration and overall survival were similar with Fludara and CAP. Within the stipulated treatment period of 6 months the number of deaths was 9 (Fludara) vs. 4 (CAP).

Post-hoc analyses using only data of up to 6 months after start of treatment revealed a difference between survival curves of Fludara and CAP in favour of CAP in the subgroup of pretreated Binet stage C patients.

5.2 Pharmacokinetic properties

Plasma and urinary pharmacokinetics of fludarabine (2F-ara-A)

The pharmacokinetics of fludarabine (2F-ara-A) have been studied after intravenous administration by rapid bolus injection and short-term infusion as well as following continuous infusion and after peroral dosing of fludarabine phosphate (Fludara, 2F-ara-AMP).

No clear correlation was found between 2F-ara-A pharmacokinetics and treatment efficacy in cancer patients.

However, occurrence of neutropenia and haematocrit changes indicated that the cytotoxicity of fludarabine phosphate depresses the haematopoiesis in a dose-dependent manner.

Distribution and metabolism

2F-ara-AMP is a water-soluble prodrug of fludarabine (2F-ara-A), which is rapidly and quantitatively dephosphorylated in the human organism to the nucleoside fludarabine (2F-ara-A).

Another metabolite, 2F-ara-hypoxanthine, which represents the major metabolite in the dog, was observed in humans only to a minor extent.

After single dose infusion of 25 mg 2F-ara-AMP per m² to CLL patients for 30 minutes 2F-ara-A reached mean maximum concentrations in the plasma of 3.5 - 3.7 µM at the end of the infusion. Corresponding 2F-ara-A levels after the fifth dose showed a moderate accumulation with mean maximum levels of 4.4 - 4.8 µM at the end of infusion. During a 5-day treatment schedule 2F-ara-A plasma trough levels increased by a factor of about 2. An accumulation of 2F-ara-A over several treatment cycles can be excluded. Postmaximum levels decayed in three disposition phases with an initial half-life of approximately 5 minutes, an intermediate half-life of 1 - 2 hours and a terminal half-life of approximately 20 hours.

An interstudy comparison of 2F-ara-A pharmacokinetics resulted in a mean total plasma clearance (CL) of 79 ± 40 ml/min/m² (2.2 ± 1.2 ml/min/kg) and a mean volume of distribution (V_{ss}) of 83 ± 55 l/m² (2.4 ± 1.6 l/kg). Data showed a high interindividual variability. After intravenous and peroral administration of fludarabine phosphate, plasma levels of 2F-ara-A and areas under the plasma level time curves increased linearly with the dose, whereas half-lives, plasma clearance and volumes of distribution remained constant independent of the dose indicating a dose linear behaviour.

After peroral fludarabine phosphate doses, maximum 2F-ara-A plasma levels reached approximately 20 - 30 % of corresponding intravenous levels at the end of infusion and occurred 1 - 2 hours postdose. The mean systemic 2F-ara-A availability was in the range of 50 - 65 % following single and repeated doses and was similar after ingestion of a solution or immediate release tablet formulation. After peroral dose of 2F-ara-AMP with concomitant food intake a slight increase (<10 %) of systemic availability (AUC), a slight decrease of maximum plasma levels (C_{max}) of 2F-ara-A and a delayed time of occurrence of C_{max} was observed; terminal half-lives were unaffected.

Elimination

2F-ara-A elimination is largely by renal excretion. 40 to 60 % of the administered intravenous dose was excreted in the urine. Mass balance studies in laboratory animals with ³H-2F-ara-AMP showed a complete recovery of radio-labelled substances in the urine.

Characteristics in patients

Individuals with impaired renal function exhibited a reduced total body clearance, indicating the need for a dose reduction. *In vitro* investigations with human plasma proteins revealed no pronounced tendency of 2F-ara-A protein binding.

Cellular pharmacokinetics of fludarabine triphosphate

2F-ara-A is actively transported into leukaemic cells, whereupon it is rephosphorylated to the monophosphate and subsequently to the di- and triphosphate. The triphosphate 2F-ara-ATP is the major intracellular metabolite and the only metabolite known to have cytotoxic activity. Maximum 2F-ara-ATP levels in leukaemic lymphocytes of CLL patients were observed at a median of 4 hours and exhibited a considerable variation with a median peak concentration of approximately 20 μ M. 2F-ara-ATP levels in leukaemic cells were always considerably higher than maximum 2F-ara-A levels in the plasma indicating an accumulation at the target sites. In-vitro incubation of leukaemic lymphocytes showed a linear relationship between extracellular 2F-ara-A exposure (product of 2F-ara-A concentration and duration of incubation) and intracellular 2F-ara-ATP enrichment. 2F-ara-ATP elimination from target cells showed median half-life values of 15 and 23 hours.

5.3 Preclinical safety data

Systemic toxicity

In acute toxicity studies, single doses of fludarabine phosphate produced severe intoxication symptoms or death at dosages about two orders of magnitude above the therapeutic dose. As expected for a cytotoxic compound, the bone marrow, lymphoid organs, gastrointestinal mucosa, kidneys and male gonads were affected. In patients, severe side effects were observed closer to the recommended therapeutic dose (factor 3 to 4) and included severe neurotoxicity partly with lethal outcome (see section 4.9).

Systemic toxicity studies following repeated administration of fludarabine phosphate showed also the expected effects on rapidly proliferating tissues above a threshold dose. The severity of morphological manifestations increased with dose levels and duration of dosing and the observed changes were generally considered to be reversible. In principle, the available experience from the therapeutic use of Fludara points to a comparable toxicological profile in humans, although additional undesirable effects such as neurotoxicity were observed in patients (see section 4.8).

Embryotoxicity

The results from intravenous animal embryotoxicity studies in rats and rabbits indicated an embryo-lethal and teratogenic potential of fludarabine phosphate as manifested in skeletal malformations, foetal weight loss and post implantation loss. In view of the small safety margin between the teratogenic doses in animals and the human therapeutic dose as well as in analogy to other antimetabolites which are assumed to interfere with the process of differentiation, the therapeutic use of Fludara is associated with a relevant risk of teratogenic effects in humans (see section 4.6).

Genotoxic potential, tumorigenicity

Fludarabine phosphate has been shown to cause DNA-damage in a sister chromatid exchange test, to induce chromosomal aberrations in an *in vitro* cytogenetic assay and to increase the rate of micronuclei in the mouse micronucleus test *in vivo*, but was negative in gene mutation assays and in the dominant lethal test in male mice. Thus, the mutagenic potential was demonstrated in somatic cells but could not be shown in germ cells.

The known activity of fludarabine phosphate at the DNA-level and the mutagenicity test results form the basis for the suspicion of a tumorigenic potential. No animal studies which directly address the question of tumorigenicity have been conducted, because the suspicion of an increased risk of second tumours due to Fludara therapy can exclusively be verified by epidemiological data.

Local tolerance

According to the results from animal experiments following intravenous administration of fludarabine phosphate, no remarkable local irritation has to be expected at the injection site. Even in case of misplaced injections, no relevant local irritation was observed after paravenous, intraarterial, and intramuscular administration of an aqueous solution containing 7.5 mg fludarabine phosphate/ml.

The similarity in nature of the observed lesions in the gastrointestinal tract after intravenous or intragastric dosing in animal experiments supports the assumption that the fludarabine phosphate induced enteritis is a systemic effect.

6. PHARMACEUTICAL PARTICULARS

6.1 List of excipients

Tablet core: Cellulose, microcrystalline
Lactose, monohydrate
Silica, colloidal anhydrous
Croscarmellose sodium
Magnesium stearate

Film-coat: Hypromellose
Talc
Titanium dioxide (E171)
Ferric oxide pigment, yellow (E172)
Ferric oxide pigment, red (E172)

6.2 Incompatibilities

Not applicable.

6.3 Shelf-life

3 years.

6.4 Special precautions for storage

Store in the original package to protect from moisture.
Do not store above 25°C. Do not refrigerate

6.5 Nature and contents of container

Blisters of 5 tablets each, comprising polyamide/aluminium/polypropylene thermoformable foil with a lidding foil of aluminium. The blisters are packed in a polyethylene tablet container with a child-resistant polypropylene screw cap.

Pack sizes: 15 or 20 film-coated tablets per tablet container.

Not all pack sizes may be marketed.

6.6 Special precautions for disposal

Handling and disposal

Fludara should not be handled by pregnant staff.

Procedures for proper handling should be followed according to local requirements for cytotoxic drugs. Waste material may be disposed of by incineration.

Any unused medicinal product or waste material should be disposed of in accordance with local requirements.

7. MARKETING AUTHORISATION HOLDER

Sanofi B.V.
Paasheuvelweg 25
1105 BP Amsterdam
Netherlands

8. MARKETING AUTHORISATION NUMBER(S)

PA 611/4/2

9. DATE OF FIRST AUTHORISATION/RENEWAL OF THE AUTHORISATION

Date of first authorisation: 24 October 2000
Date of last renewal: 11 October 2010

10. DATE OF REVISION OF THE TEXT

June 2024