Keep this information with you and share it with other healthcare professionals involved in your medical care or treatment.

Your Name:

_____

Doctor's Name (who prescribed Olumiant):

_____

Doctor's phone number:

_____

Patient Alert Card

Information for Patients about OLUMIANT® (baricitinib)

This document contains important information you should be aware of before and during treatment with Olumiant.

CMAT-53293 October 2026
V5 HPRA approval 06-Oct-2026

Lilly
A MEDICINE COMPANY

Pregnancy

- Do not take Olumiant if you are pregnant or suspect you may be pregnant
- Use effective contraception while taking Olumiant (and for 1 week after, if you stop treatment)
- Tell your doctor immediately if you become (or wish to become) pregnant

Infections

Olumiant may make an existing infection worse or increase the chance of you getting a new infection or increase the chance of viral reactivation. If you have diabetes or are older than 65, you may have an increased chance of getting infections. The infection can become serious if not treated. Inform your doctor immediately if you get symptoms of infection, such as:

- Fever, wounds, feeling more tired than usual, or dental problems
- A cough that won't go away, night sweats, and weight loss. These could be symptoms of tuberculosis (an infectious disease of the lungs)
- A painful skin rash with blisters. This could be a sign of a herpes zoster infection

Non-melanoma skin cancer

Non-melanoma skin cancer has been observed in patients taking Olumiant. If new skin lesions appear during or after therapy or if existing lesions change appearance, tell your doctor.

Blood clots

Olumiant may cause a condition in which a blood clot forms in your leg that may travel to your lungs. Inform your doctor immediately if you experience any of the following symptoms:

- Swelling or pain in one leg or arm
- Warmth or redness in one leg or arm
- Shortness of breath which is unexpected
- Rapid breathing
- Chest pain

Heart attack or stroke

Inform your doctor immediately if you experience any of the following:

- Severe chest pain or tightness (that may spread to arms, jaw, neck, back)
- Shortness of breath
- Cold sweat
- One-sided weakness in arm and/or leg
- Slurred speech

You can help by reporting any side effects you may get. Reporting forms and information can be found at www.hpra.ie

By reporting side effects you can help provide more information on the safety of this medicine.

Side effects should also be reported to Lilly Ireland. Please call Lilly Ireland on 01 664 0446.