What should I do if I get TLS symptoms?

If you experience any TLS symptoms during your treatment, you should:

Call your doctor or seek medical attention immediately, and

Stop taking Venclyxto®

Read the Patient Information Leaflet (PIL) for more information.

How can I reduce my risk of getting TLS?

Drink at least **1.5 to 2 litres of water daily:**
- starting from 2 days before your first dose, and
- on the day of your first dose, and
- every day during first 5 weeks of treatment.

Drinking plenty of water will help to remove high levels of blood salts through your urine.

Attend your scheduled **blood test appointments**, as they will help your doctor identify TLS.

Before your treatment with Venclyxto®, start another medicine that helps prevent the build-up of the blood salt (uric acid) in your body, as instructed by your doctor.

Important Contact Details

My name:

My emergency contact (name and phone number):

Doctor's name – who prescribed Venclyxto®:

Doctor's contact phone number:

Version number: IE-VNCLY-230006
Date of preparation: Oct/2023
Date of HPRA approval: 18-Dec-2023

abbvie

Important Safety Information for Healthcare Professionals about Venclyxto® and TLS

This patient is taking Venclyxto® for CLL which may cause Tumour Lysis Syndrome (TLS) in the first 5 weeks of treatment.

- Evaluate blood chemistries: potassium, uric acid, phosphorus, calcium and creatinine.
- Urgently correct blood chemistry abnormalities. **This can be fatal, if left untreated.**
- If you suspect TLS, contact prescribing doctor listed on this card immediately.

Please consult the Summary of Product Characteristics (SmPC) or contact AbbVie at +353 1 4287900 for more information and if you need additional copies of the patient card.

Patient Card

Safety Information about Venclyxto® (venetoclax) for Patients with Chronic Lymphocytic Leukaemia (CLL)

Venclyxto is used to treat patients with CLL.
This card contains information you need to know about Tumour Lysis Syndrome and Venclyxto®.

- Keep this patient card with you all the time.
- Show it to any doctor involved in your medical care or if you go to the emergency department.
- Always take Venclyxto® exactly as your doctor has told you to.

What should I know about TLS?

During your treatment, you may develop high levels of certain body salts (such as potassium and uric acid) in the blood. This can lead to a condition called Tumour Lysis Syndrome (TLS) within the first 5 weeks of treatment.

TLS symptoms:

- fever or chills
- nausea or vomiting
- feeling confused
- feeling short of breath
- irregular heart beat
- dark or cloudy urine
- feeling unusually tired
- muscle pain or joint pain
- fits or seizures
- abdominal pain and bloating